Nuclear Physics News International

Volume 21, Issue 2
April–June 2011

FEATURING:

Ab initio No Core Shell Model • Microscopic Evolution
of Nuclear Equilibrium Shapes

10619127(2011)21(2)

Taylor & Francis
Taylor & Francis Group

Nuclear Physics News is published on behalf of the Nuclear Physics European Collaboration Committee (NuPECC), an Expert Committee of the European Science Foundation, with colleagues from Europe, America, and Asia.

Editor: Gabriele-Elisabeth Körner

Editorial Board

Maria José García Borge, *Madrid (Chair)*

Tullio Bressani, *Torino*

Rick Casten, *Yale*

Reiner Krücken, *Vancouver*

Jan Kvasil, *Prague and EPS/NPB*

Douglas MacGregor, *Glasgow and EPS/NPB*

Hideyuki Sakai, *Tokyo*

Hans Ströher, *Jülich*

James Symons, *Berkeley*

Marcel Toulemonde, *Caen*

Editorial Office: Physikdepartment, E12, Technische Universität München,
85748 Garching, Germany, Tel: +49 89 2891 2293, +49 172 89 15011, Fax: +49 89 2891 2298,
E-mail: sissy.koerner@ph.tum.de

Correspondents

Argentina: O. Civitaressé, *La Plata*; **Australia:** A. W. Thomas, *Adelaide*; **Austria:** H. Leeb, *Vienna*; **Belgium:** G. Neyens, *Leuven*; **Brasil:** M. Hussein, *São Paulo*; **Bulgaria:** D. Balabanski, *Sofia*; **Canada:** J. Dilling, *TRIUMF*; K. Sharma, *Manitoba*; C. Svensson, *Guelph*; **China:** W. Zhan, *Lanzhou*; **Croatia:** R. Čaplar, *Zagreb*; **Czech Republic:** J. Kvasil, *Prague*; **Slovak Republic:** P. Povinec, *Bratislava*; **Denmark:** K. Riisager, *Århus*; **Finland:** M. Leino, *Jyväskylä*; **France:** G. De France, *GANIL Caen*; M. MacCormick, *IPN Orsay*; **Germany:** K. Langanke, *GSI Darmstadt*; U. Wiedner, *Bochum*; **Greece:** E. Mavromatis, *Athens*; **Hungary:** B. M. Nyakó, *Debrecen*; **India:** D. K. Avasthi, *New Delhi*; **Israel:** N. Auerbach, *Tel Aviv*; **Italy:** M. Ripani, *Genova*; L. Corradi, *Legnaro*; **Japan:** T. Motobayashi, *RIKEN*; **Mexico:** J. Hirsch, *Mexico DF*; **Netherlands:** G. Onderwater, *KVI Groningen*; T. Peitzmann, *Utrecht*; **Norway:** J. Vaagen, *Bergen*; **Poland:** B. Fornal, *Cracow*; **Portugal:** M. Fernanda Silva, *Sacavém*; **Romania:** V. Zamfir, *Bucharest*; **Russia:** Yu. Novikov, *St. Petersburg*; **Serbia:** S. Jokic, *Belgrade*; **South Africa:** S. Mullins, *Cape Town*; **Spain:** A. Gadea, *Valencia*; **Sweden:** J. Nyberg, *Uppsala*; **Switzerland:** K. Kirch, *PSI Villigen*; **United Kingdom:** P. Regan, *Surrey*; **USA:** D. Geesaman, *Argonne*; D. W. Higinbotham, *Jefferson Lab*; M. Thoenessen, *Michigan State Univ.*; H. G. Ritter, *Lawrence Berkeley Laboratory*; G. Miller, *Seattle*.

Nuclear Physics News ISSN 1061-9127

Advertising Manager

Maureen M. Williams, 28014 N. 123rd Lane,
Peoria, AZ 85383, USA
Tel: +1 623 544 1698
Fax: +1 623 544 1699
E-mail: mwilliams@cisaz.com

Circulation and Subscriptions

Taylor & Francis Group, LLC
325 Chestnut Street
8th Floor
Philadelphia, PA 19106, USA
Tel: +1 215 625 8900
Fax: +1 215 625 8914

Subscriptions

Nuclear Physics News is supplied free of charge to nuclear physicists from contributing countries upon request. In addition, the following subscriptions are available:

Volume 21 (2011), 4 issues

Personal: \$109 USD, £64 GBP, €87 Euro

Institution: \$911 USD, £549 GBP, €726 Euro

Copyright © 2011 Taylor & Francis Group, LLC. Reproduction without permission is prohibited.

All rights reserved. The opinions expressed in *NPN* are not necessarily those of the editors or publishers. The views expressed here do not represent the views and policies of NuPECC except where explicitly identified.

Contents

Editorial	3
Feature Article	
Ab initio No Core Shell Model by Bruce R. Barrett, Petr Navrátil, and James P. Vary.....	5
Facilities and Methods	
Direct Mass Measurements of Short-Lived Nuclides at the Storage Ring Facility in Lanzhou by Yuri A. Litvinov and Hushan Xu	13
The UNEDF Project by Richard Furnstahl.....	18
Meeting Reports	
25th International Nuclear Physics Conference by Jens Dilling	25
Zakopane Conference on Nuclear Physics, “Extremes of the Nuclear Landscape” by Bogdan Fornal	26
The International Symposium on New Faces of Atomic Nuclei by Takaharu Otsuka.....	29
Waltzing to the Nuclear Limits: A Symposium in Honor of Lee Riedinger by Ani Aprahamian, Mike Carpenter, Rick Casten, Jolie Cizewski, Daryl Hartley, Filip Kondev, Witek Nazarewicz, and Mark Riley.....	30
News and views	32
Calendar	44

<p><i>Cover illustration:</i> The Test Storage Ring TSR, presently at the Max-Planck-Institut für Kernphysik in Heidelberg-see article on page 33.</p>
--

Scholarpedia in Nuclear Physics

A recent development in our society is the access to information for the greatest number of citizens. Websites where we can find information on virtually all human activities multiply every day. However, it is known that the citizen must be exceedingly careful concerning information obtained on the Web. The challenge today is not obtaining access to information but the quality of that information; we all want information that we can trust.

Scholarpedia is an encyclopedia on the Web (<http://www.Scholarpedia.org/>) whose ambition is to offer encyclopedic articles of quality, with certified information in many scientific fields. Scholarpedia at this time is not a general encyclopedia; currently it focuses on the fields of computational neuroscience, dynamical systems, computational intelligence, physics and astrophysics, and in particular for what we are concerned with here, nuclear physics (http://www.Scholarpedia.org/article/Encyclopedia_of_Nuclear_Physics).

If quality of information is a primary objective, the goal of Scholarpedia is even more ambitious—to provide open-access encyclopedic peer-reviewed articles written by recognized experts elected by the public or invited by Scholarpedia editors. The articles are maintained and evolved through similar mechanisms as Wikipedia. Among Scholarpedia authors are 15 Nobel laureates, 4 Fields medalists, and hundreds of original inventors and discoverers. Thus Scholarpedia's articles can be

cited in the same way as any other peer-reviewed article. In that sense Scholarpedia has features commonly associated with open-access online academic journals. Each article has a “curator”—typically its author—who is responsible for its content. Any modification of the article must be approved by the curator before it appears in the final, approved version.

What makes Scholarpedia unique is that each article's content is the ultimate responsibility of one of the leading experts on the topic. The expert acts as each article's curator, vetting revisions that can be proposed by any verified user of Scholarpedia. The names of current curators are placed at the top of each article, signifying their ongoing involvement with and responsibility for that article. The name of the original author is given at the bottom of each article, and permanently stored in the Scholarpedia archive. Over time the leading role of an article will shift to other scholars, ensuring that it remains timely, accurate, and relevant. The process of curatorship makes Scholarpedia a unique project. Ernest Rutherford wrote the article “Radioactivity” for the 11th edition of the Encyclopedia Britannica in 1910. If Britannica had the feature of curatorship, the best experts of today would be competing with each other for the honor of becoming curator of Ernest Rutherford's article. The goal of Scholarpedia is to identify and convince today's Ernest Rutherfords to write encyclopedic articles on their fundamental

discoveries, so that 100 years from now the best experts will be willing to maintain and update these articles. Herein also lies the greatest difference between Scholarpedia and traditional print media: although the initial authorship and review are similar to a print journal, articles are not frozen and outdated, but dynamic, subject to an ongoing process of improvement moderated by their curators. This allows Scholarpedia to be up-to-date, yet maintain the highest quality of content. Upon approval, articles in Scholarpedia are archived in a journal <http://www.Scholarpedia.org/article/Special:Journal> (ISSN 1941 6016) so that they may be cited in the same way as any other peer-reviewed article.

The citation, found below the article's title, always refers to the latest approved version of the article that is shown to visitors by default. Any particular approved revision of the article can also be cited. Taking as an example an article by Prof. Alexandra Gade and Prof. C. Konrad Gelbke *The NSCL laboratory and the FRIB facility*, on the NSCL laboratory <http://www.nscl.msu.edu> and the FRIB facility <http://www.frib.msu.edu>, its different versions after approval will form an historical witness of the evolution of the NSCL laboratory and in particular of the evolution of the FRIB project. Written for the nuclear physics section of Scholarpedia, it can be cited as: Alexandra Gade and C. Konrad Gelbke (2010), Scholarpedia, 5(1):9651.

The views expressed here do not represent the views and policies of NuPECC except where explicitly identified.

When we were asked eighteen months ago to initiate the Scholarpedia section on nuclear physics we immediately realized the importance of the task both scientifically and in terms of personal commitment. Given the challenges of our discipline we both agreed to commit ourselves to this project in order to help to further popularize scientific achievements in nuclear physics and in its applications. The aim is to offer high-level scientific articles that can be read and understood by the entire scientific community, science students, science journalists, and the educated public but that may also be of interest to our colleagues. To ensure these requirements, the ideal Scholarpedia article is written in “Scientific American” or slightly more advanced style, as appropriate at least for advanced undergraduate students in that area or graduate students in adjacent areas. The role of the encyclopedia in nuclear physics is also to be involved in major scientific events, to inform journalists and put them in contact with the authors of the relevant articles.

The work was initiated with the support of some close collaborators. Today in the nuclear physics section of Scholarpedia one can find articles either accepted by the referees or still in the referring process or articles that are “scheduled,” that is, the authors have accepted to write the article but they have not yet submitted their draft. The quality of the already published articles and the enthusiasm of the first contributors strengthen our determination to bring to a successful conclusion this project. Of course much remains to be done. For this initiative to succeed, it needs all of us, all of our community to contribute and to popularize the project. This is the ultimate meaning of this article, certainly to present the nuclear physics section of Scholarpedia, to describe its objectives, but also to encourage the whole of the nuclear physics community to adopt this tool by making suggestions, proposals for authors and to arrive—why not?—at what is the ultimate goal of Scholarpedia, the authors of each article being recognized by the entire community.

NICOLAS ALAMANOS
*Institut de Recherche sur les lois
Fondamentales de l'Univers (IRFU),
CEA Saclay, France*

PAUL-HENRI HEENEN
Free University of Brussels, Belgium

Ab initio No Core Shell Model

BRUCE R. BARRETT

Department of Physics, University of Arizona, Tucson, Arizona, USA

PETR NAVRÁTIL

Theory Group, TRIUMF, Vancouver, Canada

Physics Division, Lawrence Livermore National Laboratory, Livermore, California, USA¹

JAMES P. VARY

Department of Physics and Astronomy, Iowa State University, Ames, Iowa, USA

Introduction

A long-standing goal of nuclear theory is to determine the properties of atomic nuclei based on the fundamental interactions among the protons and neutrons (i.e., nucleons). By adopting nucleon-nucleon (NN), three-nucleon (NNN), and higher-nucleon interactions determined from either meson exchange theory or QCD, with couplings fixed by few-body systems, we preserve the predictive power of nuclear theory. This foundation enables tests of nature's fundamental symmetries and offers new vistas for the full range of complex nuclear phenomena.

Basic questions that drive our quest for a microscopic predictive theory of nuclear phenomena include:

1. What controls nuclear saturation?
2. How the nuclear shell model emerges from the underlying theory?
3. What are the properties of nuclei with extreme neutron/proton ratios?
4. Can we predict useful cross-sections that cannot be measured?
5. Can nuclei provide precision tests of the fundamental laws of nature?
6. Under what conditions do we need QCD to describe nuclear structure?

among others.

Along with other *ab initio* nuclear theory groups, we have pursued these questions [1] with meson-theoretical NN interactions, such as CD-Bonn [2] and Argonne V18 [3], that were tuned to provide high-quality descriptions of the NN scattering phase shifts and deuteron properties. We

then add meson-theoretic NNN interactions such as the Tucson-Melbourne [4] or Urbana IX [5] interactions.

More recently, we have adopted realistic NN and NNN interactions with ties to QCD. Chiral perturbation theory within effective field theory (χ EFT) [6] provides us with a promising bridge between QCD and hadronic systems [7]. In this approach one works consistently with systems of increasing nucleon number [8–10] and makes use of the explicit and spontaneous breaking of chiral symmetry to expand the strong interaction in terms of a dimensionless constant, the ratio of a generic small momentum divided by the chiral symmetry breaking scale taken to be about 1 GeV/c. The resulting NN and NNN interactions, characterized by the order of the expansion retained (e.g., “next-to-next-to leading order” is NNLO) [11, 12], provide a high-quality fit to the NN data and the $A = 3$ ground-state (g.s.) properties. The derivations of NN, NNN, and so on interactions within meson-exchange and χ EFT are well

Figure 1. Schematic illustration on how Lee-Suzuki (LS) similarity transformation [14] yields an H_{eff} in a finite model space P decoupled from the infinite complementary Q space.

¹Present address.

Figure 2. Illustration of how the SRG procedure [15] weakens the strong off-diagonal couplings of an NN potential in momentum space as the flow proceeds to smaller values of λ (left to right panels).

established but are not subjects of this review. Our focus is solution of the non-relativistic quantum many-body Hamiltonian that includes these interactions using our no core shell model (NCSM) formalism. In the next section we will briefly outline the NCSM formalism [1, 13] and then present applications, results, and extensions in later sections.

The *Ab Initio* NCSM Formalism

The *ab initio* NCSM employs realistic interactions, preserves all their symmetries, and treats all A nucleons equally in a basis space of Slater determinants using a single-particle basis, such as the 3D harmonic oscillator (HO) [1, 13]. From this foundation, we show how to derive the well-known standard nuclear shell model, introduced by Maria Goeppert-Mayer and Hans D. Jensen in 1949 (Nobel Prize in physics, 1963), that treats only a small number of valence nucleons outside of inert closed shells. In addition, we show that the *ab initio* NCSM combined with the Resonating Group Method (RGM) provides the foundation for microscopic solutions of nuclear reactions with full predictive power.

In the *ab initio* NCSM, we start with the translationally invariant, intrinsic Hamiltonian for all A nucleons. All terms act on relative coordinates—there are no single-particle energies. We then add the HO center-of-mass (CM) Hamiltonian to provide a mean-field potential that improves convergence. The effects of the CM interaction are easily separated and later subtracted. Since realistic NN + NNN interactions are strong at short distances we must introduce a theoretically sound renormalization procedure to render the problem solvable in a basis space with available computer resources. We adopt a renormalization procedure specified by a similarity transformation that softens

the interactions and generates effective operators for all observables while preserving all experimental quantities in the low-energy domain. The derived “effective” interactions still act among all A nucleons and preserve all the symmetries of the initial or “bare” NN + NNN interactions. There are two such renormalization procedures that we currently employ, one called the Lee-Suzuki (LS) scheme [14] and the other called the Similarity Renormalization Group (SRG) [15].

Figure 3. Convergence of the ${}^4\text{He}$ g.s. energy with the size of the HO basis. Calculations with the bare (dashed line) and the SRG evolved (solid line) χEFT NN+NNN interactions are compared. The SRG evolution parameter $\lambda = 2 \text{ fm}^{-1}$ was used (Figure 2). The dotted line denotes the extrapolated g.s. energy (-28.5 MeV), which is close to the experiment (-28.3 MeV).

For LS renormalization, the infinite HO basis space is divided into a finite model space and an infinite excluded space by the use of projection operators P and Q , respectively. Then the LS effective Hamiltonian H_{eff} is obtained by performing a similarity transformation, X , on the bare Hamiltonian, H , and imposing the decoupling condition $QXH X^{-1}P = 0$; that is, H_{eff} has no matrix elements between the P and Q spaces, as shown schematically in Figure 1. Determination of the exact X requires the solution of the full A -body problem, which is not feasible. However, the determination of the A -nucleon H_{eff} from two- or three-body matrix elements, obtained by the solution of the two- or three-body cluster problem, results in an excellent approximation. It also ensures that one recovers the original full problem, if the model space approaches the infinite space, so that the approximation is fully controlled. The LS method can be applied to arbitrary modern NN (NN + NNN) potentials in either coordinate or momentum space.

Recently, we have also adopted the SRG approach [15] for softening the NN + NNN interactions. By varying a flow parameter, one can dial down the coupling between the high-energy and low-energy parts of the NN (or NN + NNN) interaction, as illustrated in Figure 2. These SRG Hamiltonians can be solved in any model space P , which is now a simple truncation of the infinite basis SRG Hamiltonian. Figure 3 shows such results for ${}^4\text{He}$ as a function of the P -space size given in terms of $N_{\text{max}}\hbar\Omega$, the maximum HO energy of configurations included above the unperturbed g.s. configuration. The figure clearly

shows that the accelerated rate of convergence for the softer SRG interactions over the bare NN (or NN + NNN) interaction.

NCSM Applications and Results

The results of numerous *ab initio* NCSM applications not only show good convergence with regard to increasing size of the model space P but also have been able to reproduce known properties of $0p$ -shell nuclei as well as explain existing puzzles and make predictions of, as yet, unexplained nuclear phenomenon. We list some illustrative examples here.

We display in Figure 4 the natural-parity excitation spectra of four nuclei in the middle of the $0p$ -shell with both the NN and the NN + NNN effective interactions from χEFT [16]. Overall, the NNN interaction contributes significantly to improve theory in comparison with experiment. This is especially well-demonstrated in the odd mass nuclei for the lowest, few excited states. The case of the g.s. spin of ${}^{10}\text{B}$ and its sensitivity to the presence of the NNN interaction is clearly evident.

A recent calculation has determined the Gamow-Teller (GT) matrix element for the beta decay of ${}^{14}\text{C}$, including the effect of NNN forces [17]. These investigations show that the very long lifetime for ${}^{14}\text{C}$ arises from a cancellation between $0p$ -shell NN- and NNN-interaction contributions to the GT matrix element, as shown in Figure 5. These ${}^{14}\text{C}$ results were obtained in the largest basis space achieved to date with NNN interactions, $N_{\text{max}} = 8$ ($8\hbar\Omega$) or approximately one billion configurations.

Figure 4. States dominated by $0p$ -shell configurations for ${}^{10}\text{B}$, ${}^{11}\text{B}$, ${}^{12}\text{C}$, and ${}^{13}\text{C}$ calculated at $N_{\text{max}} = 6$ using $\hbar\Omega = 15$ MeV (14 MeV for ${}^{10}\text{B}$). Most of the eigenstates are isospin $T = 0$ or $1/2$, the isospin label is explicitly shown only for states with $T = 1$ or $3/2$. The excitation energy scales are in MeV (adopted from Ref [16]).

Other noteworthy results include an explanation of the very small quadrupole moment (Q) in ${}^6\text{Li}$ due to a strong cancellation between the one- and two-body contributions to Q [18] (Figure 6). Recent calculations for ${}^{12}\text{C}$ explained the measured ${}^{12}\text{CB}$ (M1) transition from the g.s. to the $(1^+, 1)$ state at 15.11 MeV and showed more than a factor of 2 enhancement arising from the NNN interaction. Neutrino elastic and inelastic cross sections on ${}^{12}\text{C}$ were shown to be similarly sensitive to the NNN interaction and their contributions significantly improve agreement with experiment [19]. Working in collaboration with experimentalists, we uncovered a puzzle in the GT-excited state strengths in $A = 14$ nuclei [20]. Its resolution may lie in the role of intruder-state admixtures, but this will require further work.

Extensions of the NCSM for Treating Heavier Mass Nuclei

The basic idea of the *ab initio* Shell Model with a Core [21] is to use the well-established *ab initio* NCSM to solve

Figure 5. Contributions to the ${}^{14}\text{C}$ beta decay matrix element as a function of HO shell when the nuclear structure is described by the χEFT interaction (adopted from Ref. [17]). Top panel displays the contributions with (two right bars of each triplet) and without (leftmost bar of each triplet) the NNN force at $N_{\text{max}} = 8$. Contributions are summed within each shell to yield a total for that shell. The bottom panel displays the running sum of the GT contributions over the shells. Note the order-of-magnitude suppression of the 0p-shell contributions arising from the NNN force.

Figure 6. The quadrupole moment (Q) of the g.s. for ${}^6\text{Li}$ [$1^+(T=0)$] is shown in terms of one- and two-body contributions, as a function of increasing model-space size. The one- and two-body contributions and total Q are depicted as white, gray and black histograms, respectively [18].

for the core and one- and two-body terms that are needed for performing standard Shell Model (SSM) calculations for nuclei in the sd- and pf-shells. Such SSM calculations can be performed in vastly smaller model spaces than those required for converged NCSM calculations.

For illustration let us consider 0p-shell nuclei. We first perform a standard *ab initio* NCSM calculation to obtain converged eigenenergies and eigenfunctions for the $A = 6$ system (e.g., ${}^6\text{Li}$). Next, we carry out a unitary transformation of these ${}^6\text{Li}$ results into the smaller model space of $0\hbar\Omega$ excitations, which is equivalent to a neutron and a proton in the 0p-shell and the other four nucleons energetically frozen in the 0s-shell. Thus, we obtain only two-body matrix elements in the 0p-shell, although we started with a full solution of the $A = 6$ system in the NCSM approach. However, these two-body matrix elements contain all the physics of the six-nucleon system. These two-body matrix elements can be separated into a core and one- and two-body components suitable for SSM calculations. In a similar manner we can calculate the seven-body cluster in the 0p-shell by performing an *ab initio* NCSM calculation for ${}^7\text{Li}$ and transforming this result into the 0p-shell. In this case, we can also determine the three-body term in the 0p-shell. These core and one-, two-, and three-body terms can then be used to perform SSM calculations for all the nuclei in the 0p-shell, in much smaller model spaces. Effects of neglected four-body interactions appear to be small.

Figure 7. Calculated $p\text{-}^4\text{He}$ differential cross section (bottom panels) and analyzing power (top panels) for proton laboratory energies $E_p = 12, 14.32$ and 17 MeV compared to experimental data from Refs. [29, 30, 31, 32]. The SRG- $N^3\text{LO}$ NN potential with $\lambda = 2.02\text{ fm}^{-1}$ was used.

The same approach, outlined previously for obtaining the effective components of the shell-model Hamiltonian in a single major shell, for example, the $0p$ -shell, can also be utilized for computing the effective components of any physical operator in the same major shell. See Ref. [18] for details. The results of such calculations for the quadrupole moment (Q) of ^6Li are illustrated in Figure 6. Clearly, the very small Q -moment for ^6Li arises from complex many-body correlations among all six nucleons, leading to a large cancellation between the one- and two-body contributions.

The above $0p$ -shell results encourage us to extend this approach to nuclei in the sd -shell, which will require converged results for nuclei with $A = 16, 17, 18$, and 19 . In this regard we are working on a new version of the Importance Truncation approach of Roth [22] to obtain these results.

Applications to Nuclear Reactions

A realistic *ab initio* description of light nuclei with predictive power must have the capability to describe all bound and unbound states within a unified framework. *Ab initio* calculations for scattering processes involving more than four nucleons overall are challenging and still a rare exception [23]. Even calculations of resonant states are quite complicated [24]. The development of an *ab initio* theory of low-energy nuclear reactions on light nuclei is

key to further refining our understanding of the fundamental nuclear interactions among the constituent nucleons and providing, at the same time, accurate predictions of crucial reaction rates for nuclear astrophysics.

A fully *ab initio* approach to nuclear reactions based on the NCSM requires a more precise treatment of the wave-function asymptotics and the coupling to the continuum. Therefore, we have developed a new approach, the *ab initio* NCSM/RGM [25, 26], capable of simultaneously describing both bound and scattering states in light nuclei, by combining the resonating-group method (RGM) [27] with the *ab initio* NCSM. The RGM is a microscopic cluster technique based on the use of A-nucleon Hamiltonians, with fully anti-symmetric many-body wave functions built assuming that the nucleons are grouped into clusters. By combining the NCSM with the RGM, we complement the ability of the RGM to deal with scattering and reactions with the utilization of realistic interactions and a consistent microscopic description of the nucleonic clusters achieved via *ab initio* NCSM, while preserving important symmetries, including the Pauli exclusion principle and translational invariance.

Using the aforementioned NCSM/RGM formalism, we performed extensive nucleon- ^4He calculations with the SRG-evolved NN potentials. The agreement of our calculated $n\text{-}^4\text{He}$ and $p\text{-}^4\text{He}$ phase shifts with the experimental ones is quite reasonable for the S-wave, D-wave, and $^2P_{1/2}$ -wave. The $^2P_{3/2}$ resonance is positioned at higher energy in the calculation and the corresponding phase shifts are underestimated with respect to the experimental results, although the disagreement becomes less and less pronounced beyond the resonance energy. The observed

Figure 8. Calculated inelastic $^7\text{Be}(p,p')^7\text{Be}(1/2^-)$ cross section with indicated positions of the P -wave resonances (left). Calculated S -factor of the $^3\text{He}(d,p)^4\text{He}$ fusion reaction compared to experimental data (right). Energies are in the center of mass. The SRG- $N^3\text{LO}$ NN potential with $\lambda = 1.85\text{ fm}^{-1}$ ($\lambda = 1.5\text{ fm}^{-1}$) was used, respectively.

difference is largely due to a reduction in spin-orbit strength caused by the neglect of the NNN interaction in our calculations. More details are given in Ref. [28]. For energies beyond the $^2P_{3/2}$ resonance, our calculations compare favorably with the experimental data. This is shown in Figure 7, where the NCSM/RGM p - ^4He results are compared to various experimental data sets [29–32] in the energy range $E_p \sim 12$ –17 MeV.

The $^7\text{Be}(p,\gamma)^8\text{B}$ capture reaction plays a very important role in nuclear astrophysics as it serves as an input for understanding the solar neutrino flux [33]. The extrapolation of the S-factor (i.e., the cross-section divided by the Gamow factor) to astrophysically relevant energies relies on nuclear theory. We performed NCSM/RGM calculations of the p - ^7Be scattering as a necessary preparatory step to investigate the $^7\text{Be}(p,\gamma)^8\text{B}$ capture reaction [28]. In the calculation presented in Figure 8 that included the g.s. and the lowest four excited states of ^7Be we found a 2^+ state bound by 0.16 MeV corresponding to the ^8B ground state. In experiment, ^8B is bound by 137 keV [34]. The calculated lowest 1^+ resonance appears at about 0.7 MeV. It corresponds to the experimental ^8B 1^+ state at $E_x = 0.77$ MeV. This resonance dominates the inelastic cross-section as seen in the left part of Figure 8. We find a 0^+ , another 1^+ , and two 2^+ resonances not included in the current ^8B evaluation [34]. We note, however, that in the very recent Ref. [35], the authors claim the observation of low-lying 0^+ and 2^+ resonances, based on an R-matrix analysis of their p - ^7Be scattering experiment. Effects of the 0^+ , the second 1^+ , and the second 2^+ states are visible in the inelastic cross-section above the first 1^+ state resonance. On the other hand, the 3^+ resonance affects, in particular, the elastic cross section. Calculations of the $^7\text{Be}(p,\gamma)^8\text{B}$ capture within the NCSM/RGM are in progress.

The deuterium-tritium reaction is important for possible future fusion energy generation. Even though it has been well studied experimentally, its first principles theoretical understanding is important. The $^3\text{H}(d,n)^4\text{He}$ and its mirror reaction $^3\text{He}(d,p)^4\text{He}$ are also of interest for understanding primordial nucleosynthesis. In addition, the $^3\text{He}(d,p)^4\text{He}$ is one of the few reactions to present strong electron screening effects. The first *ab initio* calculations for these reactions within the NCSM/RGM framework are underway. Our first results were obtained with the SRG NN interaction with $\lambda = 1.5 \text{ fm}^{-1}$, for which the resonance energies are close to experimental values [36]. The astrophysical S-factor for the $^3\text{He}(d,p)^4\text{He}$ reaction from beam-

target experiments is compared to NCSM/RGM calculations for bare nuclei in the right panel of Figure 8. We observe a slightly different shape of the peak than that suggested by the “Trojan-horse” data from Ref. [37]. Also, no low-energy enhancement is present in the theoretical results contrary to the beam-target data of Ref. [38] affected by the electron screening.

Summary and Outlook

The *ab initio* NCSM treats all A nucleons equally with modern NN + NNN interactions and successfully describes properties of nuclei throughout the $0p$ -shell. In combination with the RGM, it provides a truly microscopic approach for nuclear reactions. Several investigations are underway to extend the *ab initio* NCSM to nuclei with $A > 16$ and to more completely unify the original *ab initio* NCSM with the NCSM/RGM approach. The outlook includes, but is not limited to:

1. Development of effective NN, NNN, and even NNNN interactions for more detailed investigations of $0p$ - and sd -shell nuclei.
2. Development of symmetry-adapted basis spaces such as $SU(3)$ [39].
3. Extension of the NCSM calculations to sd - and pf -shell nuclei (i.e., *ab initio* SM with a core).
4. Extension of the NCSM/RGM approach to nuclear reactions with more massive projectiles and three-cluster final states
5. Coupling of the binary-cluster (A - a , a) NCSM/RGM basis and the standard A -nucleon NCSM basis to unify the original *ab initio* NCSM and NCSM/RGM approaches. This will result in an optimal and balanced description of both bound and unbound states. We name this approach *ab initio* NCSM with the continuum (NCSMC).
6. Improved extrapolation techniques for estimating converged results.
7. The development of new techniques for quantifying theoretical uncertainties.

Acknowledgments

BRB acknowledges partial support of this work by the NSF under grants PHY-0555396 and PHY0854912 and JPV acknowledges partial support from DE-FG02-87ER40371 and DE-FC02-09ER41582 (SciDAC/UNEDF). Prepared in part by LLNL under Contract DE-AC52-07NA27344. PN acknowledges support from the NSERC Grant No.

401945-2011, and LLNL LDRD Grant No. PLS-09-ERD-020, and from the LLNL Institutional Computing Grand Challenge Program.

References

1. P. Navrátil, J. P. Vary, and B. R. Barrett, *Phys. Rev. Lett.* 84 (2000) 5728; *Phys. Rev. C* 62 (2000) 054311.
2. R. Machleidt, *Phys. Rev. C* 63 (2001) 024001.
3. R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, *Phys. Rev. C* 51 (1995) 38;
4. S. A. Coon, M. D. Scadron, P. C. McNamee, B. R. Barrett, D. W. E. Blatt, and B. H. McKellar, *Nucl. Phys. A* 317 (1979) 242; S. A. Coon and H. K. Han, *Few-Body Syst.* 30 (2001) 131.
5. B. S. Pudliner, V. R. Pandharipande, J. Carlson, and R. B. Wiringa, *Phys. Rev. Lett.* 74 (1995) 4369.
6. S. Weinberg, *Physica* 96A (1979) 327; *Phys. Lett. B* 251 (1990) 288; *Nucl. Phys. B* 363 (1991) 3.
7. V. Bernard et al., *Int. J. Mod. Phys. E* 4 (1995) 193.
8. C. Ordonez, L. Ray, and U. van Kolck, *Phys. Rev. Lett.* 72 (1994) 1982; *Phys. Rev. C* 53 (1996) 2086.
9. U. van Kolck, *Prog. Part. Nucl. Phys.* 43 (1999) 337.
10. P. F. Bedaque and U. van Kolck, *Annu. Rev. Nucl. Part. Sci.* 52 (2002) 339; E. Epelbaum, *Progr. Part. Nucl. Phys.* 57 (2006) 654.
11. E. Epelbaum, W. Glöckle, and Ulf-G. Meissner, *Nucl. Phys. A* 637 (1998) 107; 671 (2000) 295.
12. D. R. Entem and R. Machleidt, *Phys. Rev. C* 68 (2003) 041001(R).
13. P. Navrátil, S. Quaglioni, I. Stetcu, and B. R. Barrett, *J. Phys. G: Nucl. Part. Phys.* 36 No. 8 (2009) 083101.
14. K. Suzuki and S. Y. Lee, *Prog. Theor. Phys.* 64 (1980) 209; K. Suzuki and R. Okamoto, *Prog. Theor. Phys.* 92 (1994) 1045.
15. S. D. Glazek and K. G. Wilson, *Phys. Rev. D* 48 5863 (1993); F. Wegner, *Ann. Phys. (Leipzig)* 3 (1994) 77. E. D. Jurgenson, P. Navrátil, and R. J. Furnstahl, *Phys. Rev. Lett.* 103 (2009) 082501.
16. P. Navrátil, V. G. Gueorguiev, J. P. Vary, W. E. Ormand, and A. Nogga, *Phys. Rev. Lett.* 99 (2007) 042501.
17. P. Maris, J. P. Vary, P. Navrátil, W. E. Ormand, H. Nam, and D. J. Dean, *Phys. Rev. Lett.* 106 (2011) 202502.
18. A. F. Lisetskiy, M. K. G. Kruse, B. R. Barrett, P. Navrátil, I. Stetcu, and J. P. Vary, *Phys. Rev. C* 80 (2009) 024315.
19. A. C. Hayes, P. Navrátil, and J. P. Vary, *Phys. Rev. Lett.* 91 (2003) 012502; nucl-th/0305072.
20. A. Negret et al., *Phys. Rev. Lett.* 97 (2006) 062502.
21. A. Lisetskiy, B. R. Barrett, M. K. G. Kruse, P. Navrátil, I. Stetcu, and J. P. Vary, *Phys. Rev. C* 78 (2008) 044302.
22. R. Roth, *Phys. Rev. C* 79 (2009) 064324.
23. K. M. Nollett, S. C. Pieper, R. B. Wiringa, J. Carlson, and G. M. Hale, *Phys. Rev. Lett.* 99 (2007) 022502.
24. G. Hagen, D. J. Dean, M. Hjorth-Jensen, and T. Papenbrock, *Phys. Lett. B* 656 (2007) 169.
25. S. Quaglioni and P. Navrátil, *Phys. Rev. Lett.* 101 (2008) 092501.
26. S. Quaglioni and P. Navrátil, *Phys. Rev. C* 79 (2009) 044606.

BRUCE R. BARRETT

PETR NAVRÁTIL

JAMES VARY

- 27. K. Wildermuth and Y. C. Tang, *A Unied Theory of the Nucleus (Vieweg, Braunschweig, 1977)*.
- 28. P. Navrátil, R. Roth, and S. Quaglioni, *Phys. Rev. C* 82 (2010) 034609.
- 29. P. Schwandt, T. B. Clegg, and W. Haeberli, *Nucl. Phys. A* 163 (1971) 432.
- 30. K. W. Brokman, *Phys. Rev.* 108 (1957) 1000.
- 31. D. C. Dodder et al., *Phys. Rev. C* 15 (1977) 518.
- 32. R. A. Hardekopf and G. G. Holsen, *Phys. Rev. C* 15 (1977) 514.
- 33. E. Adelberger et al., *Rev. Mod. Phys.* 70 (1998) 1265.
- 34. D. R. Tilley et al., *Nuclear Physics A* 745 (2004) 155.
- 35. J. P. Mitchell et al., *Phys. Rev. C* 82 (2010) 011601(R).
- 36. P. Navrátil, S. Quaglioni, and R. Roth, arXiv:1009.3965 [nuc-th].
- 37. M. La Cognata et al., *Phys. Rev. C* 72 (2005) 065802.
- 38. M. A. Aliotta et al., *Nucl. Phys. A* 690 (2001) 790.
- 39. T. Dytrych, K.D. Sviratcheva, J.P. Draayer, C. Bahri, and J.P. Vary, *J. Phys. G.* 35 (2008) 123101.

Direct Mass Measurements of Short-Lived Nuclides at the Storage Ring Facility in Lanzhou

Introduction

Properties of unstable nuclei are indispensable for our understanding of nuclear structure and of nucleosynthesis processes in stars. Today, accurate data for nuclides lying far from the valley of beta-stability are required. However, these nuclides can generally be characterized by short half-lives and very small production rates, which makes their experimental investigation extremely difficult. Very fast and efficient methods are therefore needed.

The recent advent of experimental techniques based on projectile fragmentation, in-flight separation, and heavy-ion storage rings has enabled experiments with stored exotic nuclei at relativistic energies [1, 2]. Such measurements have proven to be very

Figure 2. Photo of the experimental cooler-storage ring CSRe in Lanzhou [7,8] (Photo: IMP, Lanzhou, China).

Figure 1. Schematic view of the high-energy part of the HIRFL-CSR facility in Lanzhou [7,8].

efficient. For instance, a single stored nucleus is often sufficient for the determination of its mass [3]. Also, a number of new isotopes have been discovered and their masses and lifetimes have been determined [4, 5]. Thus, experiments on nuclides with smallest production rates are feasible.

Storage Ring Facility in Lanzhou

Presently there are two storage ring facilities worldwide performing such experiments [6]. The first one is in operation for about two decades at the Helmholtz Centre for Heavy Ion Research GSI in Darmstadt, where the exotic nuclei are produced in projectile fragmentation or fission of primary beams accelerated by the heavy-ion synchrotron SIS. The fragments

are then separated in flight with the projectile fragment separator FRS and are injected into the experimental storage ring ESR, where their properties are investigated [1, 2].

The second storage ring facility for nuclear physics research, Cooler Storage Ring at the Heavy Ion Research Facility in Lanzhou (HIRFL-CSR), has been recently taken into operation at the Institute of Modern Physics, Chinese Academy of Sciences in Lanzhou (China) [7]. The laboratory portrait of the HIRFL-CSR facility has been presented in Nuclear Physics News in 2007 [8]. The layout of the facility is schematically illustrated in Figure 1. Its principle is very similar to the GSI facility. Here, the experimental cooler-storage ring CSRe is coupled to the heavy-ion

Figure 3. A part of the measured revolution time spectrum of $A = 2Z-1$ nuclei stored in the CSRe. Nuclides whose masses were measured directly for the first time are indicated with bold symbols [18].

synchrotron (main cooler-storage ring) CSRe via the in-flight fragment separator RIBLL2.

Exotic nuclei are produced by fragmenting relativistic primary beams accelerated by the CSRe in a production target located at the entrance of the RIBLL2. The nuclides of interest are separated in flight with RIBLL2 and are injected and stored in the CSRe. A photo of the CSRe is shown in Figure 2.

Isochronous Mass Spectrometry

The revolution times, T , or the revolution frequencies, f , of various stored ions are related (in first order) to their mass-to-charge ratios, m/q , via [1]:

$$\frac{\Delta T}{T} = -\frac{\Delta f}{f} = \frac{1}{\gamma_t^2} \frac{\Delta(m/q)}{(m/q)} - \left(1 - \frac{\gamma^2}{\gamma_t^2}\right) \frac{\Delta v}{v},$$

where γ is the relativistic Lorentz factor, γ_t the transition point of the storage ring, and $\Delta v/v$ denotes the velocity spread of the stored ions. This is the central formula for storage ring experiments. It is clear that

different nuclear species can be resolved in the ring by their revolution frequencies (revolution times), if the second term on the right-hand side containing $\Delta v/v$ vanishes or becomes negligibly small. However, the secondary beams inevitably have $\Delta v/v$ of the order of a few percent due to their production in a nuclear reaction.

Two complementary ways exist to minimize this term. The first one relies on the reduction of the velocity spread by stochastic and/or electron cooling and is the basis of the Schottky mass spectrometry (SMS) [9]. The second way is to tune the ring into a special isochronous ion-optical mode and to inject the ions at energies corresponding to $\gamma = \gamma_t$ [10]. In this case the velocity spread is compensated by the orbit lengths of the ions, i.e. the frequencies of the stored ions become independent of their velocity spread. This is the basis of isochronous mass spectrometry (IMS). The biggest advantage of the IMS is that it does not require beam cooling and is a unique technique for mass measurements of very short-lived nuclei [11]. Here we report on the recent mass measurements conducted at CSRe. So far only the IMS method has been employed. Experiments with cooled ions applying the SMS method are planned in near future.

The transition point of the CSRe is $\gamma_t = 1.395$. The fragments emerging the target at this energy are highly-ionized. For the case of nuclides addressed so far at the CSRe ($Z \leq 36$), one can safely assume that they were mostly stored as bare nuclei.

The revolution times were measured with a dedicated Time-of-Flight (ToF) detector [12]. The detector contains a $19 \mu\text{g}/\text{cm}^2$ thick, 40 mm in diameter carbon foil, which is inserted into the CSRe aperture. Each stored ion penetrates the foil at every revolution which causes the release of secondary electrons from the foil surface. The electrons are guided isochronously

by perpendicularly arranged electrostatic and magnetic fields to a set of micro-channel plates. The signals from the ToF detector are directly sampled with a fast digital oscilloscope. The recording time is typically 200 μs for each injection which corresponds to about 320 revolutions of the ions in the CSRe.

First Mass Measurements at CSRe

Pilot mass measurements have been conducted in December 2007. In that experiment, $N = Z$ fragments were produced in fragmentation of ^{36}Ar primary beams [13, 14]. The masses of ^{34}Cl and ^{30}P have been determined with a relative mass accuracy $\delta m/m \approx 8 \cdot 10^{-6}$ in excellent agreement with tabulated values from the Atomic-Mass Evaluation AME [15]. Though the achieved mass accuracy was rather poor, the isochronous ion-optical mode of the CSRe, the detection and data acquisition systems as well as data analysis methods have been successfully tested.

Figure 4. Regions in the temperature-density plane where ^{64}Ge is a waiting point (WP) with an effective lifetime longer than 50% of the β -decay lifetime. Shaded areas were deduced assuming one sigma uncertainties of the AME (gray) and the new (black) mass values for ^{65}As . The figure is from Ref. [18].

Following the first successful measurements, an attempt to access so far unknown masses has been undertaken [16]. Neutron-deficient $A = 2Z-1$ nuclei have been produced by fragmenting the enriched ^{78}Kr primary beams. Masses of three short-lived nuclides were obtained for the first time, namely for ^{63}Ge ($T_{1/2} = 97(22)$ ms), ^{65}As ($T_{1/2} = 128(16)$ ms), and ^{67}Se ($T_{1/2} = 136(12)$ ms) [17]. The counting statistics was very limited and only a moderate mass accuracy was achieved. For instance the mass excess value of ^{65}As was $ME(^{65}\text{As}) = -47428(\pm 530)$ keV. It has been realized that the resolving power of the CSRe mass spectrometry is mainly limited by long-term instabilities of the guiding magnetic fields, which cause revolution-time drifts of the entire spectrum for different injections into the CSRe.

The experiment on $A = 2Z-1$ nuclei was repeated in 2009 [18]. In a two-weeks run the data for about 700,000 stored ions have been acquired. A part

of the measured time-of-flight spectrum is shown in Figure 3. The revolution time difference to a selected reference nuclide has been considered instead of analyzing the absolute revolution times of the ions. By doing so, the effects due to the instabilities were to a large extent compensated and a resolving power of about 170,000 has been achieved. By employing the new analysis method, the masses for ^{63}Ge , ^{65}As , ^{67}Se , and ^{71}Kr ($T_{1/2} = 100(3)$ ms) [17] nuclides have been determined with a relative mass uncertainty of $\delta m/m \approx 2 \cdot 10^{-6}$. These four nuclides are indicated with bold symbols in Figure 3. It is important to note that only 309, 49, 104, and 28 particles, respectively, were accumulated for these four nuclides, which proves the high efficiency of the IMS measurements at the CSRe.

The achieved high mass accuracy for the new masses is sufficient to consider them in nuclear structure and astrophysics applications. One of such applications is the astrophysical rp-process, a sequence of proton captures

and β^+ decays proceeding near the proton drip line, which is believed to power energetic X-ray bursts [19]. The time of the rp-process and thus the duration of the X-ray burst is mainly determined by the waiting point (WP) nuclei [20]. The latter are the nuclei where the proton capture stalls and the rp-process has to proceed via the slow β^+ decay. One of the suggested WP nuclei is ^{64}Ge . To determine the extend to which ^{64}Ge is a waiting point, the knowledge of the proton separation energy of ^{65}As , $S_p(^{65}\text{As})$, is required.

S_p energies can be obtained by computing the Coulomb displacement energies (CDE) [21]. By using the new data, the S_p values for ^{63}Ge , ^{65}As , ^{67}Se , and ^{71}Kr nuclides could be determined [18]. The S_p energies for ^{63}Ge , ^{65}As , and ^{67}Se agree well with the corresponding CDE calculations. However, for the case of ^{71}Kr a 1.7σ discrepancy has been observed, which may point to the necessity to consider the deformation and shape-coexistence effects, present in this mass region, in future CDE calculations [18].

The obtained $S_p(^{65}\text{As}) = -90(85)$ keV confirmed for the first time experimentally that ^{65}As is unbound against proton emission at a 68.3% confidence level. The proton decay energy is so low that a direct proton emission from the ground state of ^{65}As should be very weak due to the Coulomb barrier. We note, that the $S_p(^{65}\text{As})$ agrees with the limit of $S_p > -250$ keV [22] based on the observed β -decay and theoretical estimates of the proton penetrability.

Using the new $S_p(^{65}\text{As})$ value, the degree to which ^{64}Ge is an rp-process WP nucleus can be determined. Figure 4 shows regions in the temperature-density plane where proton captures reduce the effective lifetime of ^{64}Ge to less than 50% of the β -decay lifetime, thus resulting in a less effective waiting point (no WP). The

Figure 5. Mass measurements conducted at the CSRe. Only nuclides with measured masses are shown. Employed primary beams are indicated with stars. GSI data are from Ref. [24].

temperatures and densities needed to bypass the ^{64}Ge waiting point are now rather well defined. Furthermore, it has been shown in Ref. [18] that in the framework of a one-zone X-ray burst model [20] the 2σ variation of the new $S_p(^{65}\text{As})$ value yields essentially identical light curves. Calculations showed that 89–90% of the reaction flow passes ^{64}Ge via proton capture indicating that ^{64}Ge is not a significant rp-process waiting point. In contrast, using the estimated $S_p(^{65}\text{As})$ from the AME, leads to a reduction of the proton capture flow to 54%.

Recent Measurements and Future Developments

The successful mass measurements program has been continued in 2011. Masses of eight short-lived $A = 2Z-2$ nuclides have been measured for the first time. These nuclides are ^{43}V ($T_{1/2} = 80\#$ ms), ^{46}Mn ($T_{1/2} = 36.2(4)$ ms), ^{47}Mn ($T_{1/2} = 100(50)$ ms), ^{49}Fe ($T_{1/2} = 64.7(3)$ ms), ^{52}Co ($T_{1/2} = 115(23)$ ms), ^{53}Ni ($T_{1/2} = 45(15)$ ms), ^{55}Cu ($T_{1/2} = 40\#$ ms), and ^{56}Cu ($T_{1/2} > 200\#$ ms) [17], where “#” denotes estimated $T_{1/2}$ values. The harvest of the CSRe mass measurements is illustrated on the chart of nuclides in Figure 5.

By employing a collimation of secondary beams at the FRS, a mass resolving power of about 200,000 has been achieved in IMS experiments at the ESR [23]. However, the transmission of exotic nuclei to the ESR was dramatically reduced. Owing to the new analysis method and to the high-rate capable ToF detector, a similar resolving power could be achieved at the CSRe without limiting the transmission. This is an obvious advantage if more rare nuclides will be addressed in future measurements.

A wide range of $\Delta(m/q)/(m/q) \approx 13\%$ can be covered simultaneously in one ion-optical setting of CSRe. However, the isochronous conditions are strictly fulfilled only in a small m/q -range [23]

and only small “isochronous” parts of revolution time spectra have been analyzed so far. The same effect has been observed also in the ESR experiments [23]. It is planned to install two time-of-flight detectors in one of the straight sections of the CSRe (Figure 1). This will enable in-ring measurements of velocities of individual stored ions. This additional information can be used to correct for the non-isochronous effects and to maintain the high resolving power over the entire spectrum.

In conclusion, the first mass measurements at the CSRe have delivered precision data on short-lived neutron-deficient nuclides. A measurement of a revolution time requires a few ten revolutions of the ion in the ring, that is nuclei with half-lives as short as a few ten microseconds can be studied. Furthermore, only a few ten particles are sufficient to determine the mass with a relative mass accuracy of a few 10^{-6} . Thus, the IMS at CSRe is an extremely fast and a very efficient method.

Acknowledgments

This article is entirely based on the common effort of our colleagues within the broad international collaboration. To all of them we are deeply obliged.

References

1. B. Franzke, H. Geissel, and G. Münzenberg, *Mass Spectrom. Rev.* 27 (2008) 428.
2. Yu. A. Litvinov and F. Bosch, *Rep. Prog. Phys.* 74 (2011) 016301.
3. L. Chen et al., *Phys. Rev. Lett.* 102 (2009) 122503.
4. F. Bosch et al., *Int. J. Mass Spectrom.* 251 (2006) 212.
5. L. Chen et al., *Phys. Lett.* B691 (2010) 234.
6. Yu. A. Litvinov et al., *Acta Phys. Polonica* B41 (2010) 511.
7. J. W. Xia et al., *Nucl. Instr. Meth. Phys. Res.* A488 (2002) 11.
8. W. L. Zhan, *Nucl. Phys. News* 17 (2007) 5.
9. T. Radon et al., *Nucl. Phys.* A677 (2000) 75.

10. M. Hausmann et al., *Hyperfine Interactions* 132 (2001) 289.
11. B. Sun et al., *Phys. Lett.* B688 (2010) 294.
12. B. Mei et al., *Nucl. Instr. Meth. Phys. Res.* A624 (2010) 109.
13. M. Wang et al., *Int. J. Mod. Phys.* E18 (2009) 352.
14. X. L. Tu et al., *Chin. Phys.* C33 (2009) 516.
15. A. H. Wapstra, G. Audi, and C. Thibault, *Nucl. Phys.* A729 (2003) 129.

YURI A. LITVINOV
GSI Helmholtzzentrum für
Schwerionenforschung, Darmstadt,
Germany
Max-Planck-Institut für Kernphysik,
Heidelberg, Germany
Institute of Modern Physics, Chinese
Academy of Sciences, Lanzhou, China

HUSHAN XU
Institute of Modern Physics,
Chinese Academy of Sciences,
Lanzhou, China

16. H.S. Xu et al., *Chin. Sci. Bull.* 54 (2009) 4749.
17. National Nuclear Data Center, <http://www.nndc.bnl.gov/>
18. X. L. Tu et al., *Phys. Rev. Lett.* 106 (2011) 112501.
19. R. Wallace and S. Woosley, *Astr. J. Suppl. Ser.* 45 (1981) 389.
20. H. Schatz et al., *Phys. Rev. Lett.* 86 (2001) 3471.
21. B. Brown et al., *Phys. Rev. C* 65 (2002) 045802.
22. J. A. Winger et al., *Phys. Rev. C* 48 (1993) 3097.
23. H. Geissel et al., *Hyperfine Interactions* 173 (2006) 49.
24. J. Stadlmann et al., *Phys. Lett. B* 586 (2004) 27.

The UNEDF Project

Overview

A new era has dawned for nuclear structure and reaction theory. Renewed interest in the physics of nuclei is fueled by experiments at rare isotope beam facilities, which explore new regions of exotic nuclei; by astrophysical observations and simulations, which require controlled extrapolations of the nuclear equation of state; by the use of nuclei in experiments on fundamental symmetries, which rely on robust nuclear structure information; and by nuclear energy and security needs, which include predictions of reaction cross-sections and fission fragment properties. The UNEDF collaboration of nuclear theorists, applied mathematicians, and

computer scientists is addressing this broad spectrum of physics through a comprehensive study of all nuclei, based on the most accurate knowledge of the strong nuclear interaction, the most reliable theoretical approaches, the most advanced algorithms, and extensive computational resources.

UNEDF stands for “Universal Nuclear Energy Density Functional.” The mission of this five-year SciDAC (“Scientific Discovery through Advanced Computing”) project is three-fold [1]:

- First, to find an optimal energy density functional (EDF) using all our knowledge of the nucleonic

Hamiltonian and basic nuclear properties.

- Second, to apply the EDF theory and its extensions to validate the functional using all the available relevant nuclear structure and reaction data.
- Third, to apply the validated theory to properties of interest that cannot be measured, in particular the properties needed for reaction theory.

The long-term vision is to arrive at a comprehensive and quantitative description of nuclei and their reactions. UNEDF is striving to replace phenomenological models with a well-founded microscopic theory that delivers maximum predictive power with quantified uncertainties.

By nuclear theory standards, UNEDF is large: it involves over 50 researchers from 9 universities and 7 national laboratories. Annually, it provides training to about 30 young researchers (postdocs and students). UNEDF is like a large experimental collaboration with multiple facets to its research goals.

The SciDAC model (<http://www.scidac.gov>) has led UNEDF members to form partnerships in which applied mathematicians and computer scientists work collaboratively with physicists to develop the required advanced algorithms and tools. At the same time, the project has catalyzed new cross-cutting physics collaborations, which include nuclear theorists around the world as international partners. The scope of the UNEDF project is large. Here we present brief snapshots that highlight the collaborative effort, the progress made, and the future

Figure 1. Table of the nuclides and the scope of theoretical methods to describe it.

prospects; more details and references can be found at the unedf.org website.

Forging New Connections

The playing field for UNEDF is the nuclear landscape of Figure 1. The sizable area marked “terra incognita,” which is populated by unstable isotopes with large neutron fractions, is of particular interest. Above the table of nuclides are shown three broad classes of theoretical methods, which are also used in atomic and condensed matter physics. The bulk of the nuclides are covered by Density Functional Theory (DFT), which provides

the theoretical underpinning and computational framework for building a nuclear EDF. By enhancing and exploiting the overlaps with *ab initio* and configuration interaction (CI) approaches, we are constructing and validating a universal EDF informed by microscopic interactions as well as experimental data.

The UNEDF components and paths between them are shown in the UNEDF strategy diagram in Figure 2. New interconnections and collaborative efforts within and across the boundaries have become indispensable. By necessity the developments have largely occurred in parallel; in

Figure 3. Carbon-12 density calculated with GFMC using the ADLB library [2].

subsequent sections we give examples of the interplay as UNEDF has worked to fulfill its mission.

Enhancing *ab initio* Structure Calculations

Direct connections of the EDF to microscopic nuclear interactions require improving the scope and accuracy of *ab initio* methods. Major advances in *ab initio* nuclear structure calculations under UNEDF have focused on improving the effective use of high performance computing—especially the efficient scaling to very large numbers of parallel processors.

A prime example of UNEDF synergy has been the development of the Asynchronous Dynamic Load Balancing (ADLB) software library by using Green’s Function Monte Carlo (GFMC) calculations as a test bed. Load balancing means ensuring that every parallel processor is used effectively; it is essential to optimal scaling. The ADLB library has enabled GFMC to run efficiently on over 100,000 core processors [2].

A major test case is the ^{12}C energy and proton density (see Figure 3), which is also a proof of principle whether a microscopic nuclear Hamiltonian calibrated to few-body nuclei can successfully predict the energy

Figure 2. UNEDF project scope and various active interconnections.

Figure 4. Theoretical (“*ab-initio*”) spectrum from NCFC compared to experimental results for ^{14}F [6].

and density distribution of larger nuclei. Unlike quantum chemistry, where the Coulomb interaction is firmly established, the input nuclear Hamiltonian must be tested and improved as needed even as the nuclear DFT itself is developed. The GFMC/ADLB combination predicts the ^{12}C experimental energy and radius to within 1%, which is a compelling validation [2]. The next targets are excited states in ^{12}C , including the famous Hoyle state.

Configuration interaction *ab initio* methods use basis expansions and require the (partial) diagonalization of extremely large matrices. These are known by various names according to details of the implementation, including No-Core Shell Model (NCSM) and No-Core Full Configuration (NCFC) methods (see accompanying article by Barrett, Navratil, and Vary). Improvements by applied mathematicians working with nuclear theorists on eigensolvers, data structures, and new combinatorial algorithms have resulted in substantial performance gains for CI codes such as MFDn, allowing efficient scaling to large numbers of processors [3].

An example of what this scaling enables is the recent prediction of the ^{14}F mass and excitation spectrum in advance of the experimental measurements, which meant solving a Hamiltonian

matrix of dimension 2 billion using 30,000 cores [4] (Figure 4). The predictions and measurement agree within the combined experimental and theoretical uncertainties (not shown). Another recent application involving many UNEDF members is an explanation of the anomalously long lifetime of ^{14}C by identifying the critical role of the three-nucleon (NNN) force in its beta decay [5]. We can look forward to many such confrontations of theory and experimental data from rare isotope facilities.

CI codes used within the UNEDF project apply both fully microscopic and phenomenological shell model approaches, and use different algorithms. But common computational issues are leading UNEDF scientists to develop the Leadership Class Configuration Interaction (LCCI) framework to unify, preserve, and disseminate valuable CI codes and resources. This will provide a user-friendly environment for researchers to download and run state-of-the-art CI codes needed for theoretical predictions as well as experimental analysis.

To make *ab initio* methods more convergent, it can be advantageous to “soften” the initial Hamiltonian by decoupling low-momentum and high-momentum parts. Renormalization group (RG) methods achieve this decoupling by evolving interactions in small steps. The resulting two-nucleon (NN) potentials are very convergent but the initial NNN interaction is also changed. Until recently a consistent NNN evolution had not been achieved. But a UNEDF collaboration successfully combined Similarity RG (SRG) evolution methods with NCSM technology. Figure 5 illustrates the improved convergence with the evolved three-body force [7]. These SRG interactions are now

being applied to larger nuclei and in reaction calculations.

The interactions used in the ^{14}C calculations and the SRG evolutions are systematically constructed (outside of UNEDF) using chiral effective field theory (EFT). UNEDF members have demonstrated that the coupled-cluster (CC) *ab initio* method can be used to accurately calculate closed-shell medium-mass nuclei such as $^{40,48}\text{C}$ with chiral EFT two-body interactions (or RG-softened versions) as well as proton halo nuclei like ^{19}F [8]. The CC formalism has been extended to include NNN forces and their inclusion in calculations of the heavier nuclei will break new barriers. A recent development is the first in-medium SRG diagonalization of closed-shell nuclei such as ^{40}C with accuracy comparable to CC results [9].

Microscopic Inputs to a Nuclear EDF

As implied by the diagram in Figure 2, multiple inputs and constraints are being used to build a universal nuclear EDF. The foundation is

Figure 5. Improved convergence in helium-4 using Similarity Renormalization Group (SRG) interactions, including an evolved NNN force [7].

provided by EDF's of the Skyrme type, whose phenomenological successes have been extensively benchmarked as part of the UNEDF project. *Ab initio* approaches can provide "control-data" to constrain more general functionals as well as to directly motivate novel density and momentum dependences.

Ab initio calculations can be used to test candidate EDF's even for systems not available in the laboratory. Current functionals are least constrained in their isovector dependence, so a system of neutron drops would be ideal, but the neutrons would not be self-bound. However, DFT says that the functional should work with a theoretical external potential, which can be used to "trap" neutrons and adjust their density profiles. UNEDF calculations applying GFMC (up to 16 neutrons) and Auxiliary Field Diffusion Monte Carlo (AFDMC, up to 54 neutrons) methods to generate such control-data are being used to probe deficiencies of conventional EDF parameterizations and to improve them. In Figure 6 such a comparison for radii is shown for the original SLy4 functional and an adjusted version [10]. The neutron drop results are validated by also

Figure 6. GFMC calculations of the radii of N neutrons in harmonic oscillator (HO) and Woods-Saxon (WS) traps, which are used to adjust the SLy4 Skyrme functional [10].

calculating with NCFC and with other Hamiltonians.

A parallel effort uses trapped neutron systems as well, but tests *ab initio* DFT methods against full calculations based on the same Hamiltonian. A hybrid approach uses chiral EFT for the long-distance (pion range) NN and NNN interactions that can be translated into novel density dependences through a revitalized incarnation of the density matrix expansion. These new functionals are being optimized and tested using the extensively developed EDF infrastructure [11].

Enhancing and Extending DFT Infrastructure

Much effort in the UNEDF project has been devoted to developing and improving the algorithmic and computational infrastructure needed to optimize candidate EDF's and to apply symmetry restoration required for accurate calculations of self-bound nuclei. Performance optimization of DFT-solver codes has enabled large-scale mass table calculations on 9,060 processors for 840,000 configurations in 9,000 nuclei in a 12-hour run. The MADNESS (Multiresolution ADaptive NumERical Scientific Simulation) framework is an example of applying state-of-the-art applied mathematics technology to create a DFT solver with an adaptive pseudo-spectral method [12].

Another example is the optimization algorithm POUNDerS, developed by a UNEDF team of applied mathematicians, which yields dramatic computational savings over alternative optimization methods, as seen in Figure 7. Using the UNEDF Experimental Database and the optimized solver HFBTHO, the Skyrme SLy4 functional was re-optimized using the

Figure 7. Greatly improved convergence is seen with the novel optimization algorithm POUNDerS [13].

derivative-free POUNDerS algorithm. The resulting parameterization UNEDF0 sets a solid baseline of nuclear ground-state properties to compare with future functionals [13]. New hybrid functionals with microscopic input from chiral EFT [11] are being optimized using this approach.

A New Era for Reaction Theory

One of the principal aims of the UNEDF project is to calculate reliable reaction cross-sections for astrophysics, nuclear energy, and national security, for which extensions of standard phenomenology is insufficient. The interplay of structure and reactions is essential for a successful description of exotic nuclei as well. Such interplay is characteristic of the *ab initio* no-core shell model/resonating-group method (NCSM/RGM), which treats bound and scattering states within a unified framework using fundamental interactions between all nucleons. A quantitative proof-of-principle calculation of this approach is shown in Figure 8 [14]. A wide range of applications is now possible including $^3\text{H}(d,n)^4\text{He}$ fusion, the $^7\text{Be}(p,\gamma)^8\text{B}$ reaction important for solar neutrino physics, and many more to come.

Figure 8. Calculated [14] NCSM/RGM cross-section for 17 MeV neutrons on ^4He compared to experiment.

Neutron reactions on heavier nuclei are being modeled using DFT results to predict not just bound states, but also scattering states for nucleons. Microscopic calculations of reaction cross-sections for nucleon-nucleus scattering have been performed by coupling the elastic channel to all particle-hole excitations in the target, and also with one-nucleon pickup channels. Target excitations were described in a random-phase (QRPA) framework using a Skyrme functional and the resulting transition potentials were used in large coupled-channel calculations. As illustrated in Figure 9, the calculated reaction cross-sections agree very well with experimental data, and also with predictions of global optical potentials where data are not available. For the first time, the observed absorption in the reaction cross-section can be accounted for by explicit channel coupling [15].

Another important capability for reactions is the calculation of level densities. A new proton-neutron algorithm for the parallel JMOMENTS code was recently designed and implemented, which improved its scalabil-

ity to tens of thousands of cores, and increased its overall performance by a factor of more than 10,000. This development opens the door to calculating accurate nuclear level densities and reaction rates for a large class of nuclei [16].

Cold Atoms as a Testing Ground

UNEDF theorists have made important contributions to the study of strongly coupled superfluid systems such as ultracold Fermi atoms, which show many similarities to the cold nuclear matter found in the crust of neutron stars. Cold atoms make excellent laboratories for testing and improving the computational methods to be used for nuclei; indeed, GFMC and AFDMC calculations from UNEDF scientists have set the standard for numerical results of the unitary gas at zero temperature.

Cold atom systems allow predictions of superfluid DFT that are testable against experiment. An example is shown in Figure 10, where a nuclear DFT code adapted to the antisymmetric superfluid local density approximation (ASLDA) [17] is applied to strongly interacting spin-imbalanced atomic gases in extremely elongated traps. Families of Larkin-Ovchinnikov (LO) states with prominent transversal oscillations of the pairing potential are predicted, as indicated by the radial alignment of nodes, coexisting with a superfluid state having a smooth pairing potential [18].

A recent major UNEDF achievement is the full implementation of the time-dependent superfluid local density approximation (TDSLDA) on a 3D spatial lattice [16]. Unlike many past approaches, matrix operations are not needed and the size of the basis set it can handle is 2–3 orders of magnitude larger than previous methods, with an implementation that can use

97% of the Jaguar supercomputer at Oak Ridge National Lab. These codes are being used to simulate the unitary gas (e.g., vortex formation) [16] and a heavy nucleus under the action of various external fields. While still exploratory, these first-time simulations of this kind for fermion superfluids serve as proof of principle for an eventual treatment of fission.

Quality Control in UNEDF

Integral to the UNEDF project is the verification of methods and codes, the estimation of uncertainties, and assessment. Methods to verify and validate include the cross-checking of different theoretical methods and codes (e.g., GFMC vs. AFDMC vs. NCFC), the use of multiple DFT solvers with benchmarking, and the confrontation of *ab initio* functionals with *ab initio* structure using the same Hamiltonian. Uncertainty quantification follows using tools for correlation analysis to estimate errors and significance. A new way to estimate theory error bars is to use multiple

Figure 9. Calculated reaction cross-sections for protons on ^{90}Zr as a function of incident energy compared to data. The solid line is the full calculation while the dashed curve is from the global optical potential [15].

RG-evolved Hamiltonians and examine the cutoff dependence of calculated observables.

The UNEDF assessment component has required the development and application of statistical tools. Particularly important for potential experiments is the analysis of experimental data significance. For example, the sensitivity of two optimized functionals to particular data is shown in Figure 11 [13]. Statistical tools are used to deliver uncertainty quantification and error analysis for theoretical studies as well as for the assessment of new experimental data. Such technologies are virtually unknown in the low-energy nuclear theory community at present, but are essential as new theories and computational tools are applied to entirely new nuclear systems and to conditions that are not accessible to experiment.

Outlook

These brief highlights represent only part of the development within the UNEDF project and new results

Figure 10. Pairing potential of polarized Fermi gas in an extremely elongated trap with different aspect ratios [18] calculated using nuclear codes developed for fission.

are appearing steadily. Please visit unedf.org to find further summaries, updates, and references. While UNEDF is in the final year of its 5-year term, its impact will be ongoing. Indeed, UNEDF has created infrastructure and interconnections that are only just beginning to be fully exploited. An important spin-off for the future has been the training of young scientists in the new developments of low-energy nuclear theory.

The worldwide impact of the UNEDF collaboration, which is unique in the field of low-energy nuclear physics in its scope, size, and structure, was evident at the 2010 INPC meeting in Vancouver, where UNEDF highlights were quoted in several plenary talks. While based and funded in the United States, UNEDF collaborates closely with foreign efforts and individual scientists sharing similar fundamental science goals. Such collaborations include joint software developments and benchmarking, and representatives of international collaborating projects attend the UNEDF annual meetings. These UNEDF activities have had positive leveraging effects on our foreign partners, including the joint UNEDF-JUSTIPEN and UNEDF-FIDIPRO efforts and a new initiative involving low-energy nuclear theory in France, FUSTIPEN.

High performance computing provides answers to questions that neither experiment nor analytic theory can address; hence, it becomes a third leg supporting the field of nuclear physics. A series of meetings on extreme scale computing in 2009 identified several major research thrusts that have a strong foundation in UNEDF achievements [20]. These include pathways to a quantitative

Figure 11. The sensitivity of candidate EDF fits to particular data can be assessed. See Ref. [13] for details.

microscopic description of fission, computing properties of nuclei that determine the r-process nucleosynthesis path in stars, computing properties of nuclei used in double-beta decay experiments and neutrino-nucleus cross-sections for modeling supernova explosions, and computing the triple-alpha process that produces ^{12}C , the nucleus at the core of organic chemistry and thus life forms. The future prospects are bright!

Acknowledgments

This work was supported by the UNEDF SciDAC Collaboration under DOE Grant DE-FC02-07ER41457.

References

1. G. Bertsch et al., *SciDAC Review* 6 (2007) 42.
2. E. Lusk et al., *SciDAC Review* 17 (2010) 30.
3. P. Maris et al., *Procedia Comp. Sci.* 1 (2010) 97.
4. P. Maris et al., *Phys. Rev. C* 81 (2010) 021301(R).
5. P. Maris et al., *Phys. Rev. Lett.* 106 (2011) 202502.
6. V. Z. Goldberg et al., *Phys. Lett. B* 692 (2010) 307.
7. E.D. Jurgenson et al., *Phys. Rev. Lett.* 103 (2009) 082501.

8. G. Hagen et al., *Phys. Rev. Lett.* 101 (2008) 092502; G. Hagen et al., *Phys. Rev. Lett.* 104 (2010) 182501.
9. K. Tsukiyama et al., *Phys. Rev. Lett.* (2011), arXiv:1006.3639.
10. S. Gandolfi et al., *Phys. Rev. Lett.* 106 (2011) 012501.
11. M. Stoitsov et al., *Phys. Rev. C* 82 (2010) 054307.
12. G. I. Fann et al., *J. Physics: Conf. Ser.* 180 (2009) 012080.
13. M. Kortelainen et al., *Phys. Rev. C* 82 (2010) 024313.
14. P. Navrátil et al., arXiv:1009.3965.
15. G. Nobre et al., *Phys. Rev. Lett.*, 105 (2010) 202502.
16. R. Senkov et al., *Phys. Rev. C* 82 (2010) 024304.
17. A. Bulgac and M. M. Forbes, *Phys. Rev. Lett.* 101 (2008) 215301.
18. J. C. Pei et al., *Phys. Rev. A* 82 (2010) 021603(R).
19. A. Bulgac and K. J. Roche, *J. Phys.: Conf. Ser.* 125 (2008) 012064; A. Bulgac et al., arXiv:1011.5999.
20. www.er.doe.gov/ascr/Misc/Grand-Challenges.html

RICHARD FURNSTAHL
(FOR THE UNEDF COUNCIL)
*Department of Physics,
Ohio State University*

25th International Nuclear Physics Conference

From 4–9 July 2010, TRIUMF, Canada’s national laboratory for particle and nuclear physics, hosted the 25th International Nuclear Physics Conference (INPC) 2010 in Vancouver, Canada. The triennial INPC is the main conference in the field and is supported by IUPAP (International Union for Pure and Applied Physics). This year it attracted over 750 delegates (including 150 graduate students) from 43 countries and covered topics in nuclear structure, reactions, and astrophysics; hadronic structure, hadrons in nuclei, and hot and dense QCD; new accelerators and underground nuclear physics facilities; neutrinos and nuclei; and applications and interdisciplinary research.

The conference opened its scientific plenary program with a talk by P. Braun-Munzinger (GSI/EMMI Darmstadt) who highlighted progress since the last conference in Tokyo 2007, showcasing theoretical and experimental examples from around the world. All fields were well represented by the plenary

program and a well-attended afternoon parallel program, with over 250 invited and contributed talks and over 380 posters presented. The poster sessions were among the most lively sessions with high participation rates by graduate students and post-doctoral fellows.

Conferences like the INPC span an entire discipline, fostering many unexpected links, discussions, or collaborations among participants. Scientific highlights included presentations in the fields of Hot and Dense QCD reporting on experimental and theoretical progress at the RHIC facility. The Nuclear Reaction session provided highlights from many new and exciting facilities including the RIKEN RIBF in Japan, and an outlook for what can be expected from FAIR in Germany and FRIB in the United States. The quest toward the “Island-of-Stability” in the Superheavy Element community is still on, and new progress was reported with the identification of element 114.

Impressive progress in the theoretical sector was presented as well, in particular new ab-initio calculation approaches. Applications of these methods and progress in the nucleon-nucleon interactions were presented in the Nuclear Structure sessions, where 3-body forces interactions are now considered state of the art. Predictions of such calculations can then be tested by experiments, as presented for example for ground state properties of exotic nuclei with laser experiments and ion trap measurements. In-beam or in-flight experiments pave the way to even more exotic isotopes where new magic numbers for the nuclear Shell Model are appearing. This will also prove relevant for Nuclear Astrophysics, where significant progress was achieved experimentally with new direct capture reactions measurements with rare isotope beams and background-suppressed facilities located in underground laboratories. Neutron star research and new modeling results of core-collapse supernovae were presented, which clearly indicated the need for neutrino interactions. Neutrinos also played a large role in other sessions, such as the New Facility session, where among other new exciting projects, progress from the deep underground facilities were presented. First beam results from long-baseline oscillation experiments showed progress in this field, and double-beta decay experiments are nearing first possible results, keenly awaited by not only the community of nuclear physicist but many others as well. The Fundamental Symmetry sessions is always one of the highlights, where progress of

Figure 1. Before the storm: the conference venue, the Chan Centre of Performing Arts at the University of British Columbia.

Figure 2. The opening plenary talk gave highlights of progress in nuclear physics, presented by P. Braun-Munzinger (GSI/EMMI) in the beautiful Chan Centre auditorium at the University of British Columbia.

Standard Model tests using atomic nuclei or nuclear physics methods are used to probe sectors complementary to large particle physics experiments, for example atomic and neutron EDM experiments. Recent progress was reported in the sector of nuclear beta decay as related to the testing of the CKM Unitarity matrix, as well as the W-mass and the Weak Mixing Angle.

Talks on the muon anomalous magnetic moment and its sensitivity for probing “New Physics” showcased the burgeoning activity in the field.

INPC 2010 made a special effort to attract many graduate students and post-doctoral fellows to the conference. For example, TRIUMF combined its traditional summer school with the U.S. National Science Foundation summer

school for nuclear physics, and offered the school directly prior to the conference. This allowed the school to recruit some of the INPC delegates as lecturers, but also gave students a broad overview to the field of nuclear physics before the conference. In addition, INPC 2010 teamed up with *Nuclear Physics A* to provide awards to the best student oral presentation and the three top poster presentations at the conference. An international panel of judges together with members from the editorial board of *Nuclear Physics A* finally decided on the following award winners among a very strong field of applicants: P. Finnlay (Guelph, Canada) for oral, and Y. J. Kim (Indiana, USA), E. Rand (Guelph, Canada), and T. Brunner (Munich, Germany) for posters.

The field clearly presented itself in a healthy and dynamic state, with many young people eagerly anticipating the advent of new experiments, theory, and facilities. At the end of the conference IUPAP announced the selection of the host of the next INPC conference: it will be held in 2013 in Florence, Italy.

JENS DILLING
TRIUMF Vancouver

Zakopane Conference on Nuclear Physics, “Extremes of the Nuclear Landscape”

The Zakopane Conferences on Nuclear Physics belong to one of the oldest series of conferences in the field of nuclear physics. Their tradition is almost half a century old and 2010 was the 45th meeting in the series.

The Conference was held from 30 August through 5 September, and had

185 participants who came to Zakopane from 50 institutions from all over the world—70 of them were Ph.D. students and young researchers. The meeting was organized jointly by the Henryk Niewodniczański Institute of Nuclear Physics of the Polish Academy of Sciences (IFJ PAN) in Kraków, the Marian

Smoluchowski Institute of Physics of the Jagiellonian University, and the Committee of Physics of the Polish Academy of Sciences. This time, the Organizing Committee was led by Adam Maj, acting as chairman, Piotr Bednarczyk—Scientific secretary and Maria Kmiecik—Managing director.

The title “Extremes of the Nuclear Landscape,” which was chosen for the Conference, indicated that the scientific program would cover a broad spectrum of forefront research activities presently ongoing in the area of nuclear physics. This ambitious enterprise was carried on by the organizers with support from the Board of Conveners that gathered distinguished colleagues with expertise in various areas of the field. Each convener, acting in close collaboration with the organizers, prepared a specific program for his/her session by inviting lecturers and selecting contributions for oral presentations. According to the tradition of the Zakopane Conferences, during the construction of the scientific program special attention was paid to offering the graduate students and young researchers enthusiastic and pedagogical overviews of the most recent research subjects in Nuclear Physics from both the theoretical and the experimental points of view.

The Conference began with the two keynote talks delivered by Walter Henning and James Vary—both speakers alluded to a unifying theme, “extremes of the nuclear landscape.”

Walter Henning presented a broad overview of recent results from the existing radioactive beam facilities and illustrated progress in the construction of the new RIB research centers. In a similar manner, but with focus on the theory aspects, James Vary concentrated on prospects of getting insights into the origins of nuclear structure based on the fundamental approach tied to QCD.

The topics signaled in Vary’s talk were discussed thoroughly during the first regular session that was convened by Witek Nazarewicz (Tennessee/Warszawa) under the title “Computing Atomic Nuclei: Frontiers of Nuclear Structure Theory.” In this session, lectures were given by Achim Schwenk (TRIUMF/Darmstadt), Morten Hjorth-Jensen (Oslo), Stuart Pittel (Delaware), Frederick Nowacki (Strasbourg), Dario Vretenar (Zagreb), and Wojtek Satuła (Warszawa).

The review of the latest experimental results in nuclear structure physics started with the heaviest nuclei—this part was led by Matti Leino (Jyväskylä). Here, the lectures, largely focused on spectroscopic and radiochemical investigations of heavy elements, were pre-

sented by Dieter Ackerman (GSI), Sergey Dmitriev (Dubna), and Paul Greenlees (Jyväskylä).

The latest achievements in the study of *proton-rich nuclei* were presented in the session organized by Mike Bentley (York) in which the invited speakers were: Bob Wadsworth (York), Dirk Rudolph (Lund), Tommi Eronen (Jyväskylä), and Bertram Blank (Bordeaux).

A wealth of information on *neutron-rich nuclei*, which had been recently accumulated in experiments using radioactive beams or new reaction product detection techniques, provided a basis for the part that was convened by Faical Azaiez (Orsay). The list of speakers included: Hiroyoshi Sakurai (RIKEN), Daniel Bazin (MSU), Didier Beaumel (Orsay), Krzysztof Rykaczewski (ORNL), Reiner Krücken (München), and Giacomo de Angelis (Legnaro).

Gamma-ray spectroscopic studies of neutron-rich nuclei, although with emphasis on the reaction mechanisms, were the main focus in the series of talks named “Nuclear Reactions and Spectroscopy with Novel Techniques.” The content of this part was

constructed by Marek Lewitowicz (GANIL) who invited Silvia Leoni (Milano), Wojtek Królas (Kraków), Navin Alahari (GANIL), and Hans-Jürgen Wollersheim (GSI).

Closely related to the aforementioned topics presented were the issues discussed in the session “Nuclear Lifetimes and Collectivity” that was led by Hans Geissel (GSI). Here, the invited lecture was given by Wolfram Korten (Saclay).

The part titled “Collective Modes,” conducted by Angela Bracco (Milano), was devoted to studies of high-lying collective excitations built on both ground and highly excited states. Of main interest here were the pygmy resonances in neutron-rich nuclei, the compression-mode giant resonances, prompt dipole gamma emission as well as the soft dipole modes and the tidal-wave mode of collective excitations; all this was presented by Takashi Nakatsukasa (RIKEN), Sunniva Siem (Oslo), Franco Camera (Milano), Umesh Garg (Notre Dame), Indranil Mazumdar (Mumbai), and Walter Reviol (St. Louis).

The “Structure of Light Nuclei and Astrophysics” session, convened by Tohru Motobayashi (RIKEN), while introducing the scientific activity on nuclear clustering also showed a broad area of research on the boundary between nuclear physics and astrophysics. The list of invited speakers included Hisashi Horiuchi (Osaka), Christian Beck (Strasbourg), David Jenkins (York), Moshe Gai (Yale), and Silvio Cherubini

(Catania). This part of the meeting ended with two talks that illustrated the construction status of the two radioactive beam facilities in Europe: Marek Lewitowicz (SPIRAL2) and Zbigniew Majka (FAIR).

A separate gathering had as the objective the studies that do not belong to the main stream of research activity in low-energy nuclear physics, although they are of interest to the broad nuclear physics audience. The content was organized by Christoph Scheidenberger (GSI) under the title “Beyond Nuclear Physics.” Invited presentations were given by Josef Pochodzalla (Mainz), Jürgen Gerl (GSI), Andrzej Rybicki (Kraków), Marek Kowalski (Kraków), Peter Thirolf (München), and Ludwik Pieńkowski (Warszawa).

A special event of the Conference was The Midnight School prepared and tutored by Jerzy Dudek (Strasbourg) under the title “Symmetries in Nature—Symmetries in Nuclei.” This happening was addressed to the very general public. A unifying theme were mysterious numerical relations observed in Nature together with the well-known symmetries established in the world of atomic nuclei.

One of the evenings was entirely devoted to the poster session. The session was a very energetic event with innumerable technical conversations, which extended far beyond the scheduled time—all of which are indicators of stimulus and strong interest in our field.

The Conference had the pleasure to host as a special invited speaker

Michał Heller, the 2008 Templeton Prize Laureate. The special guest delivered a lecture devoted to Georges Lemaître’s contribution to the foundation of modern relativistic cosmology.

Altogether, the scientific program consisted of 45 invited talks, 51 oral presentations, and the poster session with 49 contributing posters.

Zakopane Schools have a reputation of offering opportunities for intense interaction between graduate students, young researchers, and senior scientists. The Zakopane Conference 2010 gave many occasions for such interactions through the discussions during scientific sessions as well as during the traditional excursions to the most scenic places in the area of Zakopane and the Tatra Mountains. Even the weather conditions, which were not the most favorable for hiking (putting it straight, for most of the time it was raining cats and dogs), contributed very positively to the overall atmosphere of the event by keeping people together, thus enabling many long discussions to take place. It was visible that the scientific and social parts of the Conference fantastically complemented each other, thus providing yet another example of the importance of both cognitive and human aspects of Science.

The organizers hope to see you at the next Zakopane Conference in 2012 (27 August–2 September).

BOGDAN FORMAL
IFJ PAN, Kraków

The International Symposium on New Faces of Atomic Nuclei

The International Symposium on New Faces of Atomic Nuclei was held on 15–17 November 2010 in Okinawa, Japan, in order to discuss future directions of nuclear physics at the 80th birthday of Professor Akito Arima. The venue was the Okinawa Institute of Science and Technology (OIST), which has recently been constructed in the main island of Okinawa for the promotion of science and technology in Okinawa in an international atmosphere. Akito Arima is one of the co-chairs of its Board of Governors, and has contributed to create this atmosphere.

This symposium was devoted to vital discussions on future directions of nuclear physics. Atomic nuclei carry most of the mass (weight) of all materials in and on the earth, and are ultimate source of all energies causing phenomena there. The complete understanding or full usage of atomic nuclei is, however, yet to come. In other words, new faces of atomic nuclei are still emerging. Recent progress of nuclear physics indeed brings us to unexplored frontiers of science. Thus, the name “New Faces” came up again.

Such new faces have not come from scratch but from various efforts made over decades, to many of which Akito Arima has contributed substantially. The symposium started with the opening greetings by Ms. H. Shou, a former vice governor of Okinawa, representing also the OIST, and myself as the chair of the organizing committee. The first scientific session was “IBM today,” where recent developments of the Interacting Boson Model (IBM) were presented. Iachello (Yale) talked about a possible application of the IBM to double beta decay. Van Isacker (GANIL) showed a new algebraic approach to peculiar properties of $N = Z$ nuclei. Nomura (Tokyo) then reported new development for the microscopic foundation of IBM, validating it by comparison to experiments. All referred to Arima’s initiatives at the early stage of IBM studies.

In the second session, Sakai (RIKEN) introduced activities to organize nuclear physics activities in Asia, by creating Asian Nuclear Physics Association (ANPhA). This talk is quite appropriate, as Arima has been very keen in developing nuclear phys-

ics in Asia. Gales (GANIL) reviewed the new era of physics with exotic nuclei with SPIRAL2 at GANIL, followed by a talk by Suzuki (Nihon) on nuclear astrophysics.

In the afternoon, Sakurai (RIKEN) gave an overview on RIKEN’s nuclear physics program, focusing on the recently built RIBF accelerator complex, and Nazarewicz (ORNL) presented another overview from the theoretical side with emphasis on superheavy elements. Three theoretical talks by Meng (Beijing), Kamimura (RIKEN), and Kanada-En’yo (Kyoto) followed.

Tonomura (Hitachi/OIST) was one of the guest speakers, and visualized quantum phenomena observed with electrons. Asakawa (Osaka) and Braun-Munzinger (GSI/EMMI) discussed QGP theoretically and experimentally. Tamura (Tohoku) introduced the current status of J-PARC, while Shimizu (Sophia) discussed hadron resonances.

On the second day, Talmi (Weizmann), Barrett (Arizona), and Shimizu (Tokyo) discussed classic and contemporary issues of the shell model. Weidenmueller (Heidelberg) talked about nuclear excitations by strong laser pulses. We had presentations by younger generations such as Kawabata (Kyoto), Hinohara (RIKEN), and Ishii (Tokyo) from clustering to lattice QCD. Hjorth-Jensen (Oslo) summarized the current status of many-body problems of nuclear structure.

We had *Interdisciplinary Links* session then. It was started by a video greeting by Dorfan, who has been a high-energy physicist at Stanford and the president-elect of OIST. Casten (Yale) gave a remarkable introduction

International Symposium “New Faces of Atomic Nuclei”

at OIST on November 16, 2010

of nuclear physics to the general audience, and Frois (CEA) showed how physics and society can be linked on the energy issue, by showing a nice movie. Skoglund (OIST) gave a talk “Freeze the macromolecules to see them dance!”

In the second part of the *Interdisciplinary Links* session, Tsuchiya (Ryukyus), who is an oceanic biologist, explained how biological interactions work in coral reefs. Balantekin (Wisconsin) went up to the sky, talking about the role of nuclear physics in understanding the universe and the origin of elements. Hirao (ANTM) has been working on ion-beam therapy and presented impressive slides to how nuclear beams cured cancer for many patients.

We had a very lively and pleasant banquet celebrating Arima’s 80th birthday at the Intercontinental Manza Beach Resort. Many former students of Arima were present. In the same year, Arima was awarded the Order of Culture, which is only award granted by the Emperor. This was celebrated also because a

nuclear physicist was awarded the Order of Culture for the first time.

On the third day, recent developments in the nuclear shell model were reported by Brown (MSU), Ginocchio (LANL), Utsuno (JAEA), and Suzuki (Niigata). Zhao (Shanghai) is still working with Arima, and talked about their collaborative work on quantum chaos, Kajino (NAO) talked about neutrinos, Bentz (Tokai) discussed nuclear magnetic moments, which is another favorite subject of Arima. Sugawara-Tanabe (Otsu) gave the final talk on Elliot SU(3) and triaxiality. Horiuchi (RCNP) has given a warm closing remark.

We see that the range of the talk subject is quite wide, reflecting the variety of the contributions made by Arima. We believe that the participants perceived good ideas on the future of nuclear physics. The proceedings will be published from AIP as one of its conference proceedings series.

The organizers were W. Benz (Kanagawa), H. Horiuchi (Osaka), M. Oka (TIT), T. Otsuka (chair, Tokyo),

and N. Yoshinaga (Saitama). The total number of participants was 62, including 6 from the United States, 3 from France, 2 from Germany, 2 from China, and 1 each from Israel and Norway. This symposium has been supported by the OIST, the EFES program of JSPS, the Inoue Science Foundation, and the CNS of the University of Tokyo.

TAKAHARU OTSUKA
University of Tokyo

Waltzing to the Nuclear Limits: A Symposium in Honor of Lee Riedinger

A conference “Waltzing to the Nuclear Limits—A symposium in honor of Lee Riedinger,” was held at the Sea Pines resort on Hilton Head Island, South Carolina, USA, 25–27 February 2011. Attended by about 80 people from the United States and abroad, the conference featured invited talks and a poster session covering a wide range of topics in contemporary nuclear structure physics, with special emphasis on those representing extremes in nuclear structure, such as exotic nuclei far from stability

and nuclei at very high spin. The conference name has a dual connotation. It recalls Maria Goeppert Mayer’s description that nuclei are “built up like an onion in layers, with the protons and neutrons revolving around each other and spinning in orbit, like couples in a waltz around a ballroom” as well as referring to Lee Riedinger’s love of dancing (albeit not specifically waltzing).

Talks focused on a variety of subjects including broken symmetries, new collective modes, nuclei at high

and ultrahigh spins, isomers, shell structure in the Sn region, the fragility of magicity, neutron transfer studies of exotic nuclei, quantum phase transitions, moments and radii, fission product studies, lifetime measurements, the role of simulations in science, and facilities and instruments such as ATLAS, HRIBF, NSCL, GANIL/SPIRAL 2, FRIB, ELI, GREY, and AGATA. New developments in all these areas attest to the health and vitality of the field and to the advances being made due to

technological revolutions in accelerator design, instrument capability, and computational power. Together, these are launching nuclear structure physics into a new era, focusing on territories of the nuclear chart never before accessible.

Lee Riedinger's many physics interests and achievements were often cited, ranging from pioneering early work on band mixing, to a wealth of studies of high spin nuclear physics, as were his more recent interests in energy, his community service as a former chair of the DNP, advisor to Senate Majority Leader Howard Baker, his long-term role as Federal budget guru to the nuclear physics community, and his important role in the quest to provide important isotopes to the nuclear and medical communities. Many of these contributions, as well as his sporting, dancing, and other pursuits were highlighted in a friendly "roast" at the Conference Banquet attended by over a hundred people, including his wife Tina, and daughters Jen and Kara, as well as

many of Lee's former students and post-docs. About 20 of Lee's colleagues and friends offered stories and reminiscences of Lee and his family. Some of Lee's many and varied career achievements are summarized in the following couplets (slightly edited) taken from a longer poem written for the occasion:

Using Sodium Iodide even and Germanium,
 Lee studied nuclei like Samarium.
 With Joe and Noah, he studied bandmixing a ton
 And with that our eternal respect he won,
 With a landmark paper in PR 179
 That was a work ever so fine.
 In Copenhagen town at the Bohr Institute
 Lee's physics really began to take root.
 What else could be in store
 Listening to Ben and Aage Bohr?
 Bandcrossing, moments of inertia, and alignments galore
 Having seen one, he wanted many more!

Where would UT and ORNL be without Lee?

He has guided both for physics and energy,

As professor and administrator,
 Teacher, and inspired mentor.

For helping our community Lee has had a flair

From budget guru to DNP Chair.

The atmosphere of the conference was warm and friendly, with new science and old remembrances, as well as the traditional Gordon Conference-like softball game. All the attendees are eagerly awaiting Lee's hundredth birthday to have another conference.

ANI APRAHAMIAN
Notre Dame

MIKE CARPENTER
Argonne National Laboratory

RICK CASTEN
Yale University

JOLIE CIZEWSKI
Rutgers University

DARYL HARTLEY
U.S. Naval Academy

FILIP KONDEV
Argonne National Laboratory

WITEK NAZAREWICZ
University of Tennessee and Oak Ridge National Laboratory

MARK RILEY
Florida State University

Official Release of the NuPECC Long Range Plan 2010

The Nuclear Physics European Collaboration Committee (NuPECC) establishes a full review of research in all fields of nuclear physics that is published every five years. The work on the “2010 Long Range Plan” has started in the summer of 2009 and the final report has been released 9 December 2010 in an official ceremony that took place at the Royal Academy in Brussels.

The document is divided into six subdomains: hadron physics, phases of strongly interacting matter, nuclear structure and dynamics, nuclear astrophysics, fundamental interactions, and nuclear physics tools and applications. Leading scientists of each subject were asked to contribute to each chapter.

The aim of this document is to review the recent achievements, putting into evidence outstanding results, and at the same time to identify the main lines of research for the next decade and to foresee the hottest topics. Different strategies are proposed to develop medium and long-term plans, with an European perspective and put it in a global context.

The elaboration of the long range involved the whole community through two meetings organized in Frankfurt October 2009 and in Madrid in May 2010. The goal of the meetings was to invite the European

Nuclear Physics community, some 6,000 scientists, to express its view on the development of Nuclear Physics in the next decade and to bring its input for a coherent development of the field. The document was prepared

during this period by six working groups, one for each subdomain that has been identified, under the supervision of NuPECC.

The final document was released in a ceremony in Brussels under the

**EUROPEAN
SCIENCE
FOUNDATION**
SETTING SCIENCE AGENDAS FOR EUROPE

FORWARD LOOK

**Perspectives of Nuclear
Physics in Europe**

NuPECC Long Range Plan 2010

auspice of the Belgian presidency of the European Union. It was open by an official representative of the federal minister Magnette, in charge of energy and environment. Talks were given by representatives of organizations supporting and organizing research in Europe, Drs. Dr. Beatrix Vierkorn-Rudolph Chair of the European Strategy Forum on Research Infrastructures, ESFRI, Dr. Marc Heppener for ESF, and Dr. Christian Kurrer for Commission's Directorate-General for Research. The long-range

plan itself was presented by the NuPECC chairman, Dr. Gunther Rosner. A scientific view on the evolution of nuclear physics in the next decades was the subject of the last talk, given by Dr. Jean-Paul Blaizot.

A special effort was made to present the long-range plan in a form accessible to the largest possible audience. To this effect, a booklet was written by science writers under the supervision of NuPECC and a short pedagogic movie was prepared by a communication agency to present the

status and the future of nuclear physics. All these documents are accessible on the NuPECC website: <http://www.nupecc.org/index.php?display=lrp2010/main>

PAUL-HENRI HEENEN
Université Libre de Bruxelles

MARIA JOSÉ GARCIA BORGE
CSK, Madrid

NuPNET: Successful Launch of NuPNET's First Common Call for Joint Transnational Activities in Nuclear Physics

On 14 February 2011, NuPNET—the joint European network for Nuclear Physics Infrastructures—launched its first common call for joint transnational activities in three important research areas of Nuclear Physics. With a total budget of not less than 3.4 million Euros, this call is co-funded by 12 agencies responsible for the funding of Nuclear Physics in 10 European countries (<http://www.nupnet-eu.org/wps/portal/launch-of-nupnet-call.html>).

NuPNET—the ERA-Net for Nuclear Physics infrastructures (<http://www.nupnet-eu.org/>)—is a three-year project launched on 27 March 2008 and financed by the European Commission to the tune of 1.3 million Euros. Coordinated by IN2P3/CNRS, this European initiative incorporates 18 funding agencies/organizations from 14 European countries, 1 associate member and NuPECC, who acts as NuPNET's "Scientific Advisory Body."

According to the "NuPNET Report 2010" (led by I. Reinhard, PT-GSI, Germany, <http://www.nupnet-eu.org/wps/portal/nupnet-report-2010.html>), the 14 "NuPNET" countries provide more than 90% of the total funding in Nuclear Physics in Europe. NuPNET can thus be considered as being truly representative of the European map in Nuclear Physics funding and as offering the best framework conditions to reach its principal aim: *enabling the nuclear physics funding agencies to pilot joint transnational activities themselves*.

In Europe, the facilities for nuclear physics are mostly operated by laboratories financed by national funding agencies and universities. Over the next decade, a number of these will see major upgrades, including the construction of a new generation of infrastructures accepted by ESFRI as truly pan-European ventures (i.e., FAIR and SPIRAL2), which is already well under way. The nuclear physics community in Europe as well

as worldwide is fragmented when it comes to financing joint actions together and it appears that not only the funding systems of various countries and institutions are different but so are their national priorities. This is where NuPNET comes in.

Once the "NuPNET Report 2010" was done, a list of opportunities for joint activities in Nuclear Physics research was defined (led by A. Bracco, INFN, Italy), the formal and legal barriers of the national funding systems were analyzed (carried out by A. Ostapczuk, NCBiR, Poland), and a funding action plan elaborated and approved (led by N. Alamanos and B. Saghai, CEA, France).

After a survey of the existing cooperation tools (led by Ch. Stoyanov, INRNE, Bulgaria), all NuPNET member institutions were consulted at various stages to define a short-list of viable topics in view of NuPNET's Common Call. And so the next step "Selecting joint activities at a transnational level" was

achieved under the leadership of José Benlliure (MICINN, Spain).

This major step was accomplished by the representatives from *all* the 18 NuPNET member institutions/organizations during the meeting of the NuPNET Governing Council at CNRS headquarters in Paris on 15 October 2010, at which the priority research themes in Nuclear Physics in view of a common call for proposals were decided. Finally, a list of three topics was voted unanimously and the NuPNET Consortium decided to open a first common call for proposals as a pilot action, which was indeed launched in February.

The three topics of this first NuPNET Call were: (1) **R&D on new detector technologies in nuclear physics:** Gamma and neutron detection technologies based on new scintillation materials and new photo-sensors (APDs, SiPMs...). Silicon and micropatterned gas tracking detectors (GEM, Micromegas) for low and high energy applications.

Large-area diamond detectors for beam monitoring or timing. (2) **R&D on EURISOL technologies: accelerator components, targets, and ion sources.** (3) **Targeted action on nuclear structure and reactions theory.** Twelve funding agencies from 10 countries participate in this first NuPNET Call: Bulgaria (INRNE), Finland (HIP and JY), France (CEA and CNRS/IN2P3), Germany (BMBF), Italy (INFN), the Netherlands (RuG), Poland (NCBiR), Romania (IFIN-HH), Spain (MICINN), and the United Kingdom (STFC). The national contributions to a virtual common pot of this first NuPNET Call amount up to 3.4 million Euros. Proposals were submitted electronically by 22 April 2011. Eligible proposals were transmitted to evaluators, who examined them with regard to seven figures of merit (scientific relevance, strategic fit to call, international competitiveness, etc.).

On 7 July 2011 the Panel Board will meet and decide. By end of July

the results of the evaluations will be known and one expects the start of the selected joint activities by autumn 2011.

The NuPNET Consortium asked for a prolongation of the project, although no additional funding from the EU Commission is expected. Upon consultation, the EU Commission agreed to prolong the NuPNET project by nine months, until the end of November 2011. This new deadline gives NuPNET the opportunity to carry out its first call under the very best auspices. Moreover, thanks to the commitment of the participating agencies in the call, the ambition of *all* the members of the NuPNET consortium will become a reality: indeed, the first nuclear physics projects co-funded by the participating NuPNET agencies are planned to start in the fall of 2011.

SYDNEY GALÈS
AND DOROTHÉE PEITZMANN
NuPNET Co-ordination Team
IN2P3/CNRS

CologneAMS becomes Operational

In 2007 the Institute für Kernphysik (IKP) together with the Institute of Geology & Mineralogy of the University of Cologne won a national DFG contest to operate a new national centre for Accelerator Mass Spectroscopy CologneAMS (<http://www.cologne-ams.de>) [1].

It was intended to improve the experimental conditions especially for the German scientists who apply the AMS technique for their research work. It was demanded that the facility is suited for the spectrometry of all standard cosmogenic nuclides like

^{10}Be , ^{14}C , ^{26}Al , ^{26}Cl , ^{41}Ca , ^{129}I , and in addition to measure sensitively heavy ions up to ^{239}U and ^{244}Pu . The heart of the AMS setup is a 6 MV Tandatron accelerator to which a low energy mass spectrometer with a negative ion source, capable to load 200 samples, and a high energy mass spectrometer with two detector systems for the measurement of the radioactive isotopes is attached. One of these detector systems is equipped with a second foil stripper unit followed by a 120° bending magnet to allow for an effective isobar suppression [2] to reduce

overloading the ionisation chamber by isobar contaminations.

In May 2010 the accelerator was delivered to Cologne. In a spectacular action it was brought into the second basement floor of the completely renovated underground laboratory of the IKP (Figure 1). The operation of the AMS facility is completely independent from experiments performed at the 10 MV Tandem [3] which is located in the first basement floor of the accelerator area. After the assembly during the summer the AMS installation and the completely

renovated accelerator laboratory were inaugurated on 1 October in the course of a three-day celebration of the 50 years of existence of the IKP (Figure 2). These festivities included a two day symposium attended by 150 friends of the IKP, an alumni evening

Figure 1. Delivery of the 6 MV Tandatron.

with 300 participants of all ages and an open door day for the general public.

Since then the function tests of the individual components of the AMS setup could be successfully completed and first test measurements for ^{14}C and ^{36}Cl as well as for other isotopes proved that the ambitious design values could be met. After the summer vacations we expect the facility to be opened for its first users. The first projects from local groups will be devoted to geological surface investigations and astrophysical problems.

Recently an agreement for cooperation was signed between the University of Cologne and the German Research Centre for Geosciences, Potsdam (GFZ). Within the framework of this agreement the GFZ will invest 850 k€ for further developments of the AMS facility. In return this investment will be compensated by reduced charges demanded for the measurement of samples from GFZ projects.

Special laboratories for sample preparation will be built at the Institute for Geology & Mineralogy where a complete service for sample preparation will be offered for external

Figure 2. Inauguration ceremony of the new AMS installation.

users in the future. The members of CologneAMS are ready to maximize their efforts to make the new center one of the leading laboratories in the field of AMS.

References

1. A. Dewald, J. Jolie, and A. Zilges, *Nucl. Phys. News* 18(3) (2008) 26.
2. M. Klein, A. Dewald, et al., ECAART, Conference proceeding, 2010.
3. J. Jolie, H. Paetz gen. Schieck, J. Ebert, and A. Dewald, *Nucl. Phys. News* 12(1) (2002) 4.

ALFRED DEWALD AND JAN JOLIE
University of Cologne

Storage-Ring Facility at HIE-ISOLDE

Stored secondary beams enable a wide range of nuclear physics experiments as has been proven in experiments over the last two decades at the cooler-storage ring ESR in Darmstadt and since very recently also at the CSRe ring in Lanzhou [1, 2]. These facilities, however, are specialized on experiments at relativistic energies. Efforts are presently under-

taken to employ the existing storage rings also for nuclear physics experiments at lower energies, which inevitably requires the still inefficient and time consuming slowing down of stored ion beams. Therefore it is of interest to explore the possibility of installing a storage ring at an ISOL facility which naturally delivers low emittance low energy beams.

A dedicated workshop (TSR@ISOLDE) has been organized at the Max-Planck-Institute for Nuclear Physics (MPIK) in Heidelberg on 28–29 October 2010. The workshop has collected about 50 participants from all over the world. New physics ideas that can be enabled employing a low-energy storage ring at ISOLDE have been discussed. As a

Figure 1. A schematic view of the heavy-ion storage ring TSR [4]. For more details see text.

result, a Letter of Intent has been submitted to the ISOLDE and Neutron Time-of-Flight Committee, which was positively evaluated in February 2011 [3].

It is suggested to move the existing heavy-ion storage ring TSR to HIE-ISOLDE. The heavy-ion storage ring TSR is in operation at MPIK since 1988 (Figure 1). The circumference of the TSR is 55.4 m and its maximal magnetic rigidity is 1.5 Tm. Each of the four symmetric focusing periods consists of two 45° dipole magnets, five quadrupole magnets, and three sextupole magnets. A schematic view of the TSR ring is shown in the figure. The TSR has four 5.2 m long straight sections which offer ideal conditions for setting up different experiments. The electron cooler allows achieving stored beams with extremely small horizontal and vertical emittances. A dedicated ultra-cold electron target has been developed in the TSR for high-resolution experiments with electrons. Also the laser cooling can be applied to a selected number of heavy ion species. The TSR has an RF resonator

that can be used to accelerate or decelerate stored beams. A peculiar property of the TSR is a high momentum acceptance of about $\pm 3\%$, which may be used for storing exotic ions in several charge states or different radioactive nuclides with similar mass-over-charge ratios. An overview of stored beam intensities and beam lifetimes achieved in the TSR can be found on the TSR webpage [4].

New physics ideas for possible incoming experiments at HIE-ISOLDE have been suggested in the LOI. In the following we briefly sketch them:

- Experimental information on capture reactions (p, γ) and (α, γ) is very scarce and is so far restricted to stable isotopes. The stored beam is intersected by an internal gas-jet target and the recoils are measured with high efficiency after a bending magnet. The high beam intensities achieved by accumulation in the TSR can allow moving away from the stability reaching vp - and later rp -process nuclei.
- One- and two-nucleon transfer reactions, due to their selectivity, provide unique information on nuclear structure. The secondary beams from HIE-ISOLDE can be efficiently stored and cooled in the TSR. The low gas-jet target thickness is compensated by the recirculation and accumulation of ions. Owing to the excellent beam energy definition and the absence of straggling in the target, a superior energy resolution is expected.
- Atomic charge states can dramatically modify nuclear decay constants. Investigations of nuclear half-lives as a function of atomic charge states and of spin-parities

for the parent and daughter nuclei are proposed. Such experiments can be used, for instance, to address the electron screening in beta decay.

- It is predicted that 20–30% of ${}^7\text{Be}$ in the core of the Sun is present as hydrogen-like ions. However, the half-life of hydrogen-like ${}^7\text{Be}$ could not be measured up to now, which will be feasible at TSR@ISOLDE.
- Nuclear isomeric states are important probes to explore nuclear structure. Owing to the sensitivity to single stored ions, the storage-ring mass spectrometry is a unique tool to study such long-lived rare nuclear species.
- Di-electronic recombination (DR) is a well-established atomic physics research program at the TSR and the ESR. A broad scientific program can be pursued at ISOLDE, where isotopic or/and isotonic shifts can be measured thus providing information on nuclear charge radii. Furthermore, using resonant character of DR, purification of beams in ground or alternatively isomeric states is feasible, in principle. Experiments with such purified beams may be suitable for studies on laser interaction with the nuclei or for the search of predicted Nuclear Excitation by Electron Capture (NEEC) phenomenon.
- Beta beams are today one of the possible long baseline facilities to explore neutrino properties, primarily neutrino oscillation physics including CP violation in the leptonic sector. The (anti-)neutrinos are produced by acceleration and final storage of beta-decaying isotopes. The TSR@ISOLDE can be used as a versatile tool to

investigate different technical aspects. One of the key studies before a full-scale experiment is envisaged is the efficient production and storage of ${}^6\text{He}/{}^{18}\text{Ne}$ or ${}^8\text{Li}/{}^8\text{B}$ ion beams.

- Last, but not least, the high resolution of storage rings may be employed for cleaning the ion beam from isobaric contaminants.

The realization of the proposed project requires several improvements of the present REX- ISOLDE facility. These improvements include an upgrade of the present REX-EBIS to a Super-EBIT. Moreover, the TSR ring needs about $20\text{ m} \times 20\text{ m}$ area for

installation, which inevitably requires an extension of the ISOLDE experimental hall. Technical aspects of the realization of the project are being presently worked out in detail by the collaboration.

References

1. B. Franzke, H. Geissel, and G. Münzenberg, *Mass Spectrom. Rev.* 27 (2008) 428
2. Yu.A. Litvinov & F. Bosch, *Rep. Prog. Phys.* 74 (2011) 016301
3. A. Andreyev et al., “Storage ring facility at HIE-ISOLDE,” LOI to INTC, 2011.
4. Ion Storage Ring TSR, <http://www.mpi-hd.mpg.de/blaum/storage-rings/tsr/index.en.html>

KLAUS BLAUM
Max-Planck-Institut für
Kernphysik

YURI A. LITVINOV
Max-Planck-Institut für
Kernphysik
GSI Helmholtzzentrum für
Schwerionenforschung

From Proton Beam Eye Radiotherapy to a Scanning Proton Gantry—Nuclear Physics and Hadron Radiotherapy in Krakow, Poland

Figure 1. The optical bench and patient treatment chair of the proton radiotherapy facility at IFJ PAN (the beam enters from the far right toward the viewer).

On 18 February 2011 the proton radiotherapy treatment has been completed for our first two ocular patients at the Institute of Nuclear Physics of the Polish Academy of Sciences (IFJ PAN) in Kraków, Poland. This was the first time proton radiotherapy has been made clinically available to patients in Central Europe. A 38-year-old male and a 42-year female patient, both suffering from eye melanoma, underwent a series of four fractions of irradiations by a 60 MeV proton beam from the in-house-developed AIC-144 cyclotron. Nuclear physicists collaborated with physicians from the Department of Ophthalmology and Ophthalmic Oncology of the Jagiellonian University’s Collegium Medicum and

Figure 2. A preliminary architectural concept of the new National Centre for Hadron Radiotherapy in Kraków to be operational in 2014. The IBA C-235 Proteus cyclotron will produce a 230 MeV proton beam for proton radiotherapy and research in nuclear physics. The proton gantry with scanning beam will be used for advanced cancer treatment.

from the Centre of Oncology in Kraków. Thus, the clinical trial within which a total of ten patients will be treated in Kraków in 2011, has now began. The facility will next be used for regular treatment of eye-cancer patients.

The Henryk Niewodniczanski Institute of Nuclear Physics has had quite a long tradition in hadron radiotherapy. Between the years 1976 and 1994, 550 patients with cancers located in the head-and-neck or breast had been treated with a beam of fast neutrons produced on a thick beryl-

lium target using 12 MeV deuterons accelerated by the IFJ's U-120 cyclotron. This cyclotron was later replaced by the AIC-144 isochronous cyclotron in-house-designed at the IFJ PAN and adapted to proton radiotherapy between 2008 and 2010. The beam delivery system and treatment room were also in-house designed and built. Our 60 MeV proton radiotherapy facility, the only one in Poland (a 40-million country) and the first to operate in Central-Eastern Europe, is not only able to treat all Polish ocular melanoma patients (some 100 cases

per year) but also patients in neighboring European countries.

Why are protons so useful in cancer radiotherapy? Protons are able to deliver a much higher dose of ionizing radiation to the tumor volume and are able to better spare the neighboring critical organs or healthy tissues than other radiotherapy modalities. This is of particular importance to pediatric patients in whom the possible occurrence of secondary radiation-induced cancer should be minimized. Protons of initial energies ranging between 60 MeV and 250 MeV when stopping in the irradiated tissues, will deposit most of their energy (or dose) at their stopping ends, a phenomenon known as the Bragg peak. Also, unlike in photon beams, proton ranges are extremely well defined by their initial energy, thus critical organs or healthy tissues immediately behind the tumor volume can be spared by carefully adjusting the proton energy. Moreover, the entrance dose of a proton beam is lower than that of any megavolt photon beam, thus further sparing tissues upstream of the irradiated tumor volume.

The range of 60 MeV protons in water is about 30 mm, which is sufficient to treat tumors in the eye-ball but to reach deeper located tumors elsewhere in the body higher proton energies are needed, to be supplied by more powerful (and more expensive) accelerators. Poland, as a new member of the European Union is receiving some 60 billion Euro over the years 2007–2013, for reconstructing its infrastructure. Of this sum, IFJ PAN received about 50 M€ (85% from EU structural funds and 15% from Polish government) to finance the installation in Krakow of a Proteus C-235 230 MeV proton

cyclotron, produced by the Belgian Ion Beam Application (IBA) company. Construction of the new cyclotron building (National Centre of Hadron Radiotherapy—Cyclotron Centre Bronowice; Figure 2) with experimental and treatment rooms began on 17 March 2011, to be completed by the end of the year 2013.

To be able to deliver beams of these high-energy protons to any tumor location in the patient's body, a gantry is needed. The tender for an

up-to-date radiotherapy treatment room with a gantry-rotating beam will be announced within a month. The gantry is a huge construction, 11 meters in diameter and weighing 150 tons. It carries the beam-bending magnets and is able to aim the beam with sub-millimeter precision. We expect this complex, called the Cyclotron Centre in Bronowice, to begin treating patients in 2014.

Besides the proton therapy, a nuclear physics research program is

planned to be carried on. It will include among others the gamma-ray spectroscopy (giant resonances, isomer decays, etc.) and nuclear reaction (three-body forces, nuclear symmetries, fission and spallation mechanism, etc.) studies. In addition, in-beam detector tests are foreseen. We are looking forward for collaboration!

PAWEŁ OLKO
IFJ PAN, Kraków

100 Years of Nuclear Physics: The Legacy of Ernest Rutherford

Just over two years ago, I was invited to address the New Zealand High Commission in London on the centenary of Ernest Rutherford's Nobel Prize for chemistry in 1908. It was on that occasion that I met his great granddaughter, Mary Fowler, a professor of geophysics at the University of London, from whom I have since picked up several insights about the great man's life and views. This year, we celebrate what is for many scientists a far more momentous anniversary: the centenary of the publication of Rutherford's 1911 paper on the atomic nucleus in the *Philosophical Magazine* (vol. 21, p. 669).

I cannot of course do justice in this short article to his incredible legacy to science that sprang from that landmark paper. In any case, there is little doubt in my mind that in terms of his achievements, influence, leadership, and impact on science, I would rank Rutherford as the greatest experimental physicist of all time, and certainly the greatest experimental scientist of the last century. And while we nuclear

physicists rightly celebrate this year as the centenary of the paper that launched our discipline, I would like to use this opportunity to outline some of the many other key contributions Rutherford made to science.

He was born near Nelson, in New Zealand in 1871 and, after studying at Canterbury College in Christchurch, won a scholarship to come to England to work at JJ Thomson's famous Cavendish Laboratory in Cambridge. Initially, he joined "JJ" in the study of the newly discovered mysterious X-rays, but his attention was soon to drift to what would be his life-long passion: radioactivity, also newly discovered and equally mysterious.

Radioactivity

Rutherford's early research was so impressive that in 1898, when still only just 27 years old, he was offered a full professorship at McGill University in Montreal, Canada. It was there that he carried out the work that won him the Nobel Prize for a discovery

that would mark the end of chemistry's most cherished belief: that radioactivity was a process that could transform one element into another. Somehow, one type of atom emitted a particle and became another type of atom. No doubt mindful of the ill-repute arising from thousands of years of futile attempts to make gold from lead, Rutherford is reported to have said to his collaborator Frederick Soddy: "Don't call it transmutation, Soddy, or they'll have our heads as alchemists!"

By 1904, he had become the world's leading expert on radioactivity and had written the first ever textbook on the subject entitled, simply, *Radio-activity*. In it we find the famous quote: "The phenomena exhibited by the radioactive bodies are extremely complicated and some form of theory is essential to connect in an intelligible manner the mass of experimental facts that have now been accumulated."

Of course, Rutherford could not have predicted the quantum revolution

that took place in the first quarter of the 20th century. The book, which he dedicated to JJ Thomson, describes all of his work at McGill with Soddy, including his coining of alpha and beta radiation and the concept of the half-life (even if the term itself came a few years later). It was the 2nd edition of the book published in 1905 that would become *the* standard textbook on the subject, providing a complete review of what was known at the time.

The Age of the Earth

It was also in 1904 that Rutherford finally fully understood how to estimate the age of the earth: that if the radioactive elements in the Earth's crust, such as radium, thorium, actinium, and uranium, have been active since our planet was formed then their proportions now could tell us how long they had been there. Rutherford would call on two different pieces of evidence for this. First, he knew he could measure the amount of helium gas trapped in rocks. Since he was confident that his alpha particles were in fact ionized helium atoms, the amount of helium would tell him how long the radioactive materials had been decaying in the rocks and hence their age.

But the second method was the one that would capture the imagination of the scientific community at the time. The great 19th-century physicist and inventor, William Thompson (later Lord Kelvin) had worked out the age of the earth many years earlier to be one or two orders of magnitude younger than the correct value, based on a model in which it started off in a molten state and has been cooling ever since. This figure worried geologists (and biologists) who wanted a figure of many hundreds of millions

of years. However, taking into account the heat given off by radioactive elements in the earth's core meant that it had to be much older than Kelvin's estimate in order to have cooled down to its current temperature. Rutherford delivered a famous lecture at the Royal Institution in London in May 1905. Kelvin—by this time the grand old man of British science—was in the audience. Rutherford knew this and was nervous, for he knew that it would not be a wise move to publicly embarrass Kelvin by so dramatically revising his value for the age of the earth. Luckily, it seemed Kelvin was asleep during the lecture. Then, at the crucial moment, Rutherford noticed that he had opened one eye. It was at this moment that Rutherford made his famously diplomatic remark: “*Kelvin had limited age of earth* provided no new sources of heat found. *That prophetic utterance referred to what we are now considering!*” He would later say that, on hearing this, Kelvin “beamed” at him.

“Single Scattering”

And so we come to the famous alpha scattering experiment of 1909. Rutherford had moved from McGill to the University of Manchester in 1907 and it was there that he oversaw the great experiment carried out by his assistants, Hans Geiger and Ernest Marsden in 1909 in which a beam of alpha particles from a radium source were fired at a gold leaf. They observed with amazement how one in every eight thousand of these particle “bullets” bounced right back from the thin gold sheet. Of course, every physicist today knows the explanation, but it took Rutherford over a year to work out what was happening. When he did, he published it in his paper: “The scattering of alpha and

beta particles by matter and the structure of the atom.” In it, he explained the existence of the tiny dense atomic nucleus by coming up with a scattering formula—one that he worked out from the classical theory of electrostatic deflection of charge, but which can be derived quantum mechanically too (using Born Approximation). With this new model of the atom, Rutherford would overthrow the old “plum pudding” model proposed by his mentor, J J Thomson.

And so nuclear science was born, and Rutherford replaced Thomson as the dominant figure in British science.

Over half a century later, the same reasoning that led Rutherford to conclude the existence of the tiny dense nucleus at the heart of the atom would point to the quark structure of nucleons, this time via electron scattering from hydrogen (protons) in what is referred to as deep inelastic scattering.

I recall, as young research student, deriving what is known as the Schrödinger-equivalent Dirac equation containing what is known as the “Darwin term” in the nuclear potential. I did not know it at the time, but this term is named after Charles Galton Darwin (grandson of the great Darwin) who worked under Rutherford in Manchester. The young Darwin recalls an occasion around 1910 while at dinner one evening when Rutherford first talked about the idea of “single scattering” of the alpha particles and how they trace a parabolic path through the atom, much like the path of a comet passing close by the sun's gravitational field. Indeed, a copy of Newton's *Principia* was found on Rutherford's desk around this time opened to the page on the inverse square law of gravity and the bending of trajectories by

hyperbolas. And it was a source of great satisfaction to Rutherford that his classical formula proved to be correct even when derived quantum mechanically.

The First Nuclear Reaction

By 1914, Rutherford's team in Manchester was looking at how alpha particles scattered from light atoms, such as hydrogen. They realized that the hydrogen nuclei should be kicked forward by the heavier alphas. Around this time, Marsden had noticed an anomalous result in which many more hydrogen nuclei than expected were being produced in alpha scattering from a range of light targets. He wondered whether this was due to contamination in his radioactive source, or even if the source was spitting out hydrogen as well as helium nuclei. Of course, although we know now that certain heavy nuclei do indeed decay via proton emission, this was not what was happening here. In fact, what Marsden had done without realizing it was split the atom—this, in 1914! His experiment was the very first artificial nuclear reaction $^{14}\text{N}(\alpha, p)^{17}\text{O}$. On this occasion it took Rutherford not 18 months, but four years to figure out what was happening. In between 1915 and 1919, when Rutherford published four landmark papers, what was happening slowly became clearer. In the last of his 1919 papers, Rutherford wrote about his own series of nuclear reaction experiments: "We must conclude that the nitrogen atom is disintegrated under the intense forces developed in a close collision with a swift alpha particle, and that the hydrogen atom which is liberated formed a constituent part of the nitrogen nucleus."

It was also around this time that Rutherford made yet another

prophetic remark, which reflects just how pivotal his thinking was on those around him. It is well known that James Chadwick discovered the neutron in 1932. However, it was Rutherford who first suggested its possible existence. In his Bakerian lecture at the Royal Society in 1920 he writes of "... the possible the existence of an atom of mass nearly 2, carrying one charge, which is to be regarded as an isotope of hydrogen."

Here he is postulating the existence of the deuteron (the study of which would provide the material for my own Ph.D.). He then deduces from this "... the possible existence of an atom of mass 1, which has zero charge."

He even coined the term neutron for this particle.

Moonshine?

Despite making an early statement regarding the splitting of the atomic nucleus that "some fool in a laboratory might blow up the universe unawares," Rutherford was by the early 1930s dismissive of the energy locked up within it, saying "The energy produced by the breaking down of the atom is a very poor kind of thing. Anyone who expects a source of power from the transformation of these atoms is talking moonshine."

However, around the time of the discovery of the neutron in 1932 (and long before the discovery of nuclear fission) he is also said to have spoken privately to Maurice Hankey, Britain's "man of secrets," who had been Secretary of the Committee of Imperial Defence in World War I. Hankey said Rutherford told him that.. "the experiments on nuclear transformation which he was supervising in Cambridge... might one day turn out to be of great importance to the defence of the country. He did not quite know in

what way this would be so...but some inner sense, for which he apparently had no scientific justification, . . . warned him that someone should . . . 'keep an eye on the matter.'"

Further evidence of his interest in the possibility of atomic energy comes from Rutherford's interest in Fermi's experiments in Rome with slow neutrons. In the last year of his life, Bohr and Wheeler wrote their seminal paper on the liquid drop model of the nucleus. After a lecture on this topic by Bohr, he was overheard booming: "When Niels, if in a nuclear reaction mass disappears, energy will appear and, ultimately, whatever its initial form, be degraded to heat. It might be used."

Science Funding

Rutherford's main influence on science funding and policy came through the DSIR (Department of Scientific and Industrial Research), which had been founded during the Great War, in 1915. After WWI, the DSIR became the main body funding British scientific research. Today of course, it has been replaced by the U.K. Research Councils. Scientists now write grant proposals for research funding and forget that this was not always the way science has been funded. Until less than a century ago, it tended to be through private benefactors, universities, learned societies or industry; but not government—not for basic research.

Rutherford began tapping the DSIR for funds around 1924. He secured a salary for Chadwick at the Cavendish, of which he was now the director, for the post of deputy director to carry out administrative work that would free up Rutherford for more research. He also successfully applied for £8000 worth of funding

for equipment, thus marking the launch of his research in “Big Science,” the legacy of which we see today in projects from FAIR at GSI to the LHC at Cern. Rutherford would, late in life, become a very powerful figure in U.K. politics, but never ceased to devote his efforts to helping secure funding for his

researchers and promote science around the world.

Rutherford was a loud, straight-talking giant of a man who did not suffer fools gladly. He would push his colleagues, assistants, and students hard in the lab and would offer encouragement all the time. The English chemist, Henry Tizard, once

said of him: “He never considered whether he should be a grave man or a merry man, but just let inclination for the time take its course.”

JIM AL-KHALILI
University of Surrey

The French–US Theory Institute for Physics with Exotic Nuclei

A new initiative in nuclear theory has been launched: the French-US Theory Institute for Physics with Exotic Nuclei (FUSTIPEN). The Institute is based at GANIL, and is dedicated to promoting collaborative projects between researchers in the United States and in France in the area of physics of or with exotic nuclei, including nuclear structure and reaction theory, nuclear astrophysics, and tests of the standard model rare isotopes.

The Institute was inaugurated with a two-day meeting 18–19 January, attended by about 80 participants (Figure 1). The first day was devoted to the broad picture of the research

underway on exotic nuclei. Perspectives on the projects planned and underway were given by representatives of the major funding agencies, namely T. Hallman of the U.S. Department of Energy, F. Staley of the CEA/DSM, and J. Martino of the French CNRS/IN2P3. There were numerous examples showing the importance of international collaboration in the large experimental projects.

The second day was a workshop highlighting theoretical problems that could benefit from collaborative efforts, as well as reporting on ongoing French-U.S. collaborations in the physics of exotic nuclei. High on the

list of problems was the need to improve on present implementations of mean-field theory, because of its role as the foundation of nuclear structure theory. In specific regions of the nuclear chart the interacting shell model is the most powerful spectroscopy tool available for theory. The implementations of this model at the intersection of structure and reaction theory remain one of the major future challenges. There is also a great need for better reaction theory to make quantitative interpretations of experiments. Besides these key areas, there were many topics that were also presented, ranging from shell structure evolution, isospin breaking, astrophysical reaction rates, to giant resonances and fission.

The FUSTIPEN Institute will operate by funding visits of U.S.-based scientists to GANIL to collaborate on a one-to-one basis with French-based scientists. The visits can also include stays at the home institutions of the French-based researcher. Funding for FUSTIPEN is provided by the Office of Nuclear Physics of the U.S. Department of Energy while GANIL provides the local support. The French grant supplies funding for French physicists to visit

Figure 1. Inauguration of FUSTIPEN, 18–19 January 2011 in GANIL, Caen, France.

GANIL to collaborate with U.S. visitors. While FUSTIPEN is dedicated to nuclear theory, collaborations between theorists and experimentalists are essential for understanding and planning of experiments, and such activities are encouraged by the Institute. The present funding of FUSTIPEN permits the support of about 10–15 collaborative visits per year. More details about the Institute and the application process can be found on its website: fustipen@ganil.fr.

Numerous FUSTIPEN activities are already on the books. On 3 February 2011, the topical meeting “Neutron-Proton pair correlations in $N \sim Z$ nuclei” was devoted to specific correlation effects due to the isoscalar pairing. On 3 March 2011, the topical meeting “Effective field theories for nuclear structure studies” investigated current issues posed by the many-body problem of nuclear structure to effective field theories. Further activities, such as the

topical meeting on “Probing two-nucleon correlations in reactions,” and workshops on “Open quantum systems” and “Theory of nuclear fission” are planned later in this year.

G. BERTCH, S. GALES,
W. NAZAREWICZ, AND
M. PŁOSZAJCZAK
FUSTIPEN

June 13–17

München, Bavaria, Germany. XIV International Conference on Hadron Spectroscopy HADRON2011
<http://hadron2011.de/>

June 13–17

Stockholm, Sweden. Nordic Conference on Nuclear Physics 2011
<http://www.nuclear.kth.se/NCNP2011/Home.html>

June 17–20

Northampton, Massachusetts, USA. NuSYM11 International Symposium on Nuclear Symmetry Energy
<http://www.smith.edu/nusym11/>

June 27–July 2

Constanta, Romania. “Advanced many-body and statistical methods in mesoscopic systems”
<http://www.theory.nipne.ro/Constanta-Meso2011/>

June 30–July 2

RIKEN, Wako, Saitama, Japan. Joint International Symposium on Frontier of gamma-ray spectroscopy (gamma11)
<http://www.cns.s.u-tokyo.ac.jp/gamma11/>

July 4–10

St. Petersburg, Russia. Isomers in Nuclear and Interdisciplinary Research Meeting (INIR-2011)
<http://onlinereg.ru/inir2011>

July 18–22

Quito, Ecuador. IX Latin American Symposium on Nuclear Physics and Applications
<http://www.lasnpa-quito2011.org/>

August 8–12

Manchester, UK. Rutherford Centennial Conference on Nuclear Physics
<http://rutherford.iopconfs.org/>

August 28–September 2

Guelph, Canada. Fourteenth International Symposium on Capture Gamma-Ray Spectroscopy and Related Topics
<http://cgs14.physics.uoguelph.ca/>

September 5–9

Vienna, Austria. International Conference on Exotic Atoms and Related Topics - EXA2011
<http://www.oeaw.ac.at/smi/research/talks-events/exotic-atoms/exa-11/>

September 11–18

Piaski, Poland. XXXII Mazurian Lakes Conference on Physics
<http://www.mazurian.fuw.edu.pl/>

September 16–24

Erice, Sicily, Italy. Erice School 2011 “From Quarks and Gluons to Hadrons and Nuclei”
<http://crunch.ikp.physik.tu-darmstadt.de/erice/>

September 19–23

Prague, Czech Republic. Third International Workshop on Compound-Nuclear Reactions and Related Topics (CNR*11)
<http://www-ucjf.troja.mff.cuni.cz/cnr11>

October 9–14

Frascati, Italy. 8th International Conference on Nuclear Physics at Storage Rings STORI'11
<http://www.lnf.infn.it/conference/stori11/>

October 11–15

Kyoto, Japan. Yukawa International Seminar “Frontier Issues in Physics of Exotic Nuclei” (YKIS2011)
<http://www2.yukawa.kyoto-u.ac.jp/~ykis2011/ykis/index.html>

November 14–17

Wako, Japan. The 11th International Symposium on Origin of Matter and

Evolution of Galaxies (OMEG11)

<http://www.cns.s.u-tokyo.ac.jp/omeg11>

November 23–28

Hanoi, Vietnam. International Symposium on Physics of Unstable Nuclei 2011 (ISPUN11)
<http://www.inst.gov.vn/ispun11/>

November 24–25

Athens, Greece. Workshop on “Thermonuclear Reaction Rates for Astrophysics Applications”
<http://libra.inp.demokritos.gr/THERRAA/>

2012

January 23–27

Bormio, Italy. 50th International Winter Meeting on Nuclear Physics
<http://www.bormiomeeting.com/>

February 22–25

Bormio, Italy. 1st Topical Workshop on Modern Aspects in Nuclear Structure “Advances in Nuclear Structure with arrays including new scintillator detectors”
<http://www.mi.infn.it/WSBormio-Milano2012/>

March 26–29

Pisa, Italy. Direct Reactions with Exotic Beams DREB2012
<http://dreb2012.df.unipi.it/Dreb/WELCOME.html>

May 28–June 2

St. Petersburg, Florida, USA. Eleventh Conference on the Intersections of Particle and Nuclear Physics CIPANP 2012
<http://cipanp2012.triumf.ca/>

September 17–21

Bucharest, Romania. European Nuclear Physics Conference EuNPC 2012
<http://www.ifin.ro/eunpc2012/>

More information available in the Calendar of Events on the NuPECC website: <http://www.nupecc.org/>